

ANNEX B: DETAILED DATA TABLES

Section 1 data tables

Table B.1: Estimated prevalence of 12-month mental disorders, by age and sex, ADF and ABS study

Age group (Years)	Males				Females				Persons						
	ABS %	ABS 95% CI	N ADF	ADF %	95% CI	ABS %	ABS 95% CI	N ADF	ADF %	95% CI	% ABS	ABS 95% CI	N ADF	% ADF	95% CI
18-27	24.2	18.8-29.6	2,517	26.8	18.6-34.9	27.9	23.7-32.1	558	26.5	17.1-35.9	24.7	20.2-29.3	3,076	26.7	19.8-33.6
28-37	19.0	13.8-24.1	3,297	26.0	16.5-35.5	26.0	22.2-29.8	584	21.9	15.8-28.0	20.0	15.7-24.4	3,881	25.3	17.3-33.2
38-47	16.3	10.6-22.1	2,507	17.8	14.3-21.3	21.8	16.1-27.4	425	24.3	17.1-31.6	17.0	11.9-22.0	2,932	18.5	15.3-21.7
48-57	16.2	10.1-22.3	986	14.8	9.8-19.8	16.2	11.3-21.0	66	23.2	6.9-39.5	16.2	10.6-21.8	1,052	15.2	10.3-20.0

Table B.2: Estimated prevalence of 12-month affective disorders, by age and sex, ADF and ABS study

Age group (Years)	Males				Females				Persons						
	ABS %	ABS 95% CI	N ADF	ADF %	95% CI	ABS %	ABS 95% CI	N ADF	ADF %	95% CI	% ABS	ABS 95% CI	N ADF	% ADF	95% CI
18-27	5.8	2.9-8.7	1,015	10.8	5.6-15.9	7.7	5.1-10.3	190	9.0	3.1-14.9	6.1	3.6-8.5	1,204	10.5	6.1-14.8
28-37	6.4	3.4-9.3	1,565	12.3	4.8-19.9	7.5	5.4-9.6	293	11.0	6.7-15.3	6.5	4.0-9.0	1,858	12.1	5.8-18.4
38-47	5.1	2.3-7.9	998	7.1	4.9-9.3	6.9	2.6-11.1	186	10.7	6.1-15.2	5.3	2.8-7.8	1,184	7.5	5.4-9.5
48-57	4.9	2.7-7.1	456	6.8	3.8-9.9	3.8	2.2-5.4	27	9.4	0.0-19.3	4.8	2.8-6.8	482	7.0	4.0-9.9

Table B.3: Estimated prevalence of 12-month anxiety disorders, by age and sex, ADF and ABS study

Age group (years)	Males				Females				Persons						
	ABS %	ABS 95% CI	N ADF	ADF %	95% CI	ABS %	ABS 95% CI	N ADF	ADF %	95% CI	% ABS	ABS 95% CI	N ADF	% ADF	95% CI
18–27	10.6	7.0–14.1	980	10.4	6.2–14.7	19.2	16.0–22.5	455	21.6	13.1–30.1	11.9	8.9–14.8	1,435	12.5	8.6–16.3
28–37	11.3	7.6–15.0	2,518	19.8	10.3–29.3	21.8	17.6–26.1	465	17.4	11.7–23.2	12.9	9.8–16.0	2,983	19.4	11.5–27.3
38–47	12.5	7.8–17.2	1,818	12.9	10.0–15.8	18.8	13.7–23.8	293	16.8	10.4–23.2	13.2	9.1–17.3	2,111	13.3	10.6–16.0
48–57	13.3	7.5–19.2	768	11.5	7.0–16.1	14.0	9.4–18.7	58	20.4	4.8–36.0	13.4	8.0–18.7	826	11.9	7.5–16.3

Table B.4: Estimated prevalence of 12-month alcohol disorders, by age and sex, ADF and ABS study

Age group (years)	Males				Females				Persons						
	ABS %	ABS 95% CI	N ADF	ADF %	95% CI	ABS %	ABS 95% CI	N ADF	ADF %	95% CI	% ABS	ABS 95% CI	N ADF	% ADF	95% CI
18–27	15.2	10.3–20.0	1,261	13.4	7.2–19.6	9.4	6.2–12.6	45	2.1	0.0–4.4	14.3	10.2–18.4	1,306	11.3	6.3–16.4
28–37	6.5	3.3–9.8	521	4.1	2.3–5.9	2.5	0.9–4.0	66	2.5	0.0–5.1	5.9	3.2–8.7	587	3.8	2.3–5.4
38–47	4.5	2.0–7.0	565	4.0	2.1–5.9	2.0	0.4–3.6	35	2.0	0.0–4.3	4.2	2.1–6.4	600	3.8	2.1–5.5
48–57	1.8	0.3–3.3	91	1.4	0.5–2.3	2.1	0.1–4.0	7	2.3	0.0–6.5	1.8	0.4–3.2	98	1.4	0.5–2.3

Table B.5: Estimated prevalence of 12-month affective disorders, by type and sex, ADF and ABS study

	Males				Females				Persons									
	ABS %	ABS 95% CI	ADF %	ADF 95% CI	Diff (ADF–ABS)	95% CI	ABS %	ABS 95% CI	ADF %	ADF 95% CI	Diff (ADF–ABS)	95% CI	ABS %	ABS 95% CI	ADF %	ADF 95% CI	Diff (ADF–ABS)	95% CI
Depressive episodes	2.9	(2.0, 3.9)	6.0	(4.5, 7.4)	3.0	(1.3, 4.8)	4.4	(3.3, 5.6)	8.7	(6.1, 11.3)	4.3	(1.5, 7.1)	3.1	(2.3, 4.0)	6.4	(5.1, 7.7)	3.2	(1.7, 4.7)
Dysthymia	0.7	(0.1, 1.3)	1.0	(0.5, 1.4)	0.3	(–0.5, 1.0)	1.0	(0.6, 1.5)	1.6	(0.6, 2.6)	0.6	(–0.5, 1.7)	0.8	(0.2, 1.3)	1.1	(0.6, 1.5)	0.3	(–0.4, 1.0)
Bipolar affective disorder	2.5	(1.3, 3.7)	3.1	(0.8, 5.3)	0.6	(–2.0, 3.1)	2.4	(1.4, 3.4)	1.2	(0.3, 2.0)	–1.2	(–2.5, 0.1)	2.5	(1.4, 3.5)	2.8	(0.9, 4.7)	0.3	(–1.9, 2.5)
Any affective disorder	5.7	(4.2, 7.3)	9.4	(6.8, 12.0)	3.7	(0.6, 6.7)	7.3	(5.8, 8.8)	10.2	(7.5, 12.9)	2.9	(–0.2, 6.0)	5.9	(4.6, 7.3)	9.5	(7.2, 11.8)	3.6	(0.9, 6.2)

Table B.6: Estimated prevalence of 12-month anxiety disorders, by type and sex, ADF and ABS study

	Males						Females						Persons					
	ABS %	ABS 95% CI	ADF %	ADF 95% CI	Diff (ADF-ABS)	Diff 95% CI	ABS %	ABS 95% CI	ADF %	ADF 95% CI	Diff (ADF-ABS)	Diff 95% CI	ABS %	ABS 95% CI	ADF %	ADF 95% CI	Diff (ADF-ABS)	Diff 95% CI
	Panic attack	5.8	(4.4, 7.1)	6.4	(4.7, 8.0)	0.6	(-1.6, 2.8)	11.1	(9.3, 12.9)	11.6	(8.5, 14.7)	0.5	(-3.0, 4.1)	6.5	(5.2, 7.7)	7.1	(5.6, 8.6)	0.6
Panic disorder	2.5	(1.5, 3.4)	1.2	(0.8, 1.7)	-1.2	(-2.3, -0.2)	3.0	(2.1, 3.9)	2.5	(1.3, 3.8)	-0.5	(-2.0, 1.1)	2.5	(1.7, 3.4)	1.4	(1.0, 1.9)	-1.1	(-2.1, -0.2)
Agoraphobia	1.8	(1.0, 2.6)	2.7	(1.9, 3.5)	0.9	(-0.2, 2.0)	2.5	(1.8, 3.3)	1.4	(0.4, 2.4)	-1.1	(-2.4, 0.1)	1.9	(1.2, 2.6)	2.5	(1.8, 3.2)	0.6	(-0.4, 1.6)
Social phobia	4.2	(2.8, 5.6)	3.5	(2.6, 4.3)	-0.7	(-2.4, 0.9)	6.5	(5.2, 7.8)	6.2	(3.7, 8.7)	-0.3	(-3.1, 2.5)	4.5	(3.3, 5.8)	3.8	(3.0, 4.7)	-0.7	(-2.2, 0.8)
Specific phobia	-	-	5.6	(4.2, 7.0)	-	-	-	-	8.7	(6.0, 11.4)	-	-	-	-	6.0	(4.8, 7.3)	-	-
Generalised anxiety disorder	1.6	(0.9, 2.3)	1.0	(0.5, 1.4)	-0.6	(-1.5, 0.2)	3.1	(2.0, 4.1)	1.8	(0.6, 3.0)	-1.3	(-2.9, 0.3)	1.8	(1.1, 2.4)	1.1	(0.7, 1.5)	-0.7	(-1.5, 0.04)
Obsessive-compulsive disorder	1.2	(0.7, 1.8)	3.2	(0.9, 5.5)	2.0	(-0.4, 4.4)	2.9	(1.9, 3.9)	2.7	(1.1, 4.3)	-0.2	(-2.0, 1.7)	1.4	(1.0, 1.9)	3.2	(1.1, 5.2)	1.7	(-0.4, 3.8)
Post-traumatic stress disorder	4.6	(3.4, 5.8)	8.1	(5.5, 10.6)	3.5	(0.6, 6.3)	9.0	(7.5, 10.5)	10.1	(7.3, 12.8)	1.0	(-2.1, 4.2)	5.2	(4.1, 6.3)	8.3	(6.1, 10.6)	3.1	(0.6, 5.6)
Any anxiety disorder	11.5	(9.4, 13.5)	14.2	(10.9, 17.5)	2.7	(-1.2, 6.6)	19.9	(17.7, 22.1)	18.8	(15.0, 22.5)	-1.1	(-5.5, 3.2)	12.6	(10.8, 14.4)	14.8	(11.9, 17.7)	2.2	(-1.2, 5.6)

Table B.7: Estimated prevalence of 12-month alcohol disorders, by type and sex, ADF and ABS study

	Males						Females						Persons					
	ABS %	ABS 95% CI	ADF %	ADF 95% CI	Diff (ADF-ABS)	Diff 95% CI	ABS %	ABS 95% CI	ADF %	ADF 95% CI	Diff (ADF-ABS)	Diff 95% CI	ABS %	ABS 95% CI	ADF %	ADF 95% CI	Diff (ADF-ABS)	Diff 95% CI
	Alcohol harmful use	5.5	(3.9, 7.1)	3.1	(1.7, 4.5)	-2.4	(-4.6, -0.3)	3.7	(2.4, 5.1)	1.3	(0.2, 2.5)	-2.4	(-4.2, -0.7)	5.3	(3.9, 6.7)	2.8	(1.6, 4.1)	-2.4
Alcohol dependence	3.3	(2.1, 4.5)	2.6	(1.8, 3.3)	-0.7	(-2.2, 0.7)	1.4	(0.7, 2.1)	0.9	(0.2, 1.7)	-0.5	(-1.5, 0.6)	3.0	(2.0, 4.1)	2.3	(1.7, 3.0)	-0.7	(-2.0, 0.6)
Any alcohol disorder	8.8	(6.7, 10.9)	5.6	(4.1, 7.2)	-3.2	(-5.8, -0.5)	5.1	(3.7, 6.6)	2.2	(0.9, 3.6)	-2.9	(-4.9, -0.9)	8.3	(6.5, 10.1)	5.2	(3.8, 6.6)	-3.1	(-5.4, -0.8)

Table B.8: Mean number of days in the previous four weeks totally out of role for the ADF and the ABS due to health/psychological distress by 12-month ICD-10 disorder

	Males				Females				Persons					
	ABS	ABS 95% CI	ADF	ADF 95% CI	Diff (ADF-ABS)	Diff 95% CI	ABS	ABS 95% CI	ADF	ADF 95% CI	Diff (ADF-ABS)	Diff 95% CI		
Any affective disorder	2.8	(0.4, 5.3)	1.8	(1.0, 2.6)	-1.0	(-3.6, 1.5)	1.6	(0.8, 2.4)	2.1	(1.0, 3.2)	0.5	(-0.9, 1.9)	-0.8	(-3.0, 1.4)
Any anxiety disorder	2.5	(1.0, 4.0)	1.2	(0.7, 1.7)	-1.3	(-2.9, 0.3)	1.2	(0.8, 1.6)	1.6	(0.9, 2.3)	0.4	(-0.4, 1.2)	-1.0	(-2.4, 0.3)
Any alcohol disorder	0.9	(0.3, 1.5)	0.5	(0.1, 0.8)	-0.4	(-1.1, 0.3)	1.0	(0.5, 1.5)	2.7	(0.7, 4.6)	1.7	(-0.4, 3.7)	-0.3	(-0.9, 0.3)
Any disorder	1.6	(0.9, 2.4)	1.2	(0.8, 1.6)	-0.4	(-1.2, 0.5)	1.2	(0.8, 1.5)	1.4	(0.8, 1.9)	0.2	(-0.5, 0.8)	-0.3	(-1.0, 0.4)

Table B.9: Mean number of days in the previous four weeks partially out of role for the ADF and the ABS due to health/psychological distress by 12-month ICD-10 disorder

	Males				Females				Persons					
	ABS	ABS 95% CI	ADF	ADF 95% CI	Diff (ADF-ABS)	Diff 95% CI	ABS	ABS 95% CI	ADF	ADF 95% CI	Diff (ADF-ABS)	Diff 95% CI		
Any affective disorder	2.1	(1.1, 3.1)	3.0	(1.9, 4.0)	0.9	(-0.6, 2.4)	4.1	(2.7, 5.4)	6.7	(4.2, 9.1)	2.6	(-0.2, 5.4)	1.2	(-0.2, 2.6)
Any anxiety disorder	2.0	(1.3, 2.7)	2.6	(1.8, 3.4)	0.6	(-0.4, 1.7)	2.3	(1.7, 3.0)	4.2	(2.8, 5.6)	1.9	(0.3, 3.4)	0.9	(-0.1, 1.8)
Any alcohol disorder	1.6	(0.4, 2.7)	1.7	(0.8, 2.5)	0.1	(-1.3, 1.5)	1.5	(0.6, 2.4)	4.3	(0.9, 7.7)	2.8	(-0.7, 6.3)	0.3	(-1.0, 1.6)
Any disorder	1.6	(1.1, 2.1)	2.3	(1.8, 2.9)	0.8	(-0.01, 1.5)	2.3	(1.7, 2.8)	4.2	(2.9, 5.4)	1.9	(0.5, 3.3)	0.9	(0.2, 1.7)

Table B.10: Days totally unable to carry out their work, study or day-to-day activities in the previous four weeks, by type of 12-month affective disorder, subgrouped by weeks

ICD-10 affective disorder	0 days			1-7 days			8-14 days			15-21 days			22-28 days			Missing		
	N	%	95% CI	N	%	95% CI	N	%	95% CI	N	%	95% CI	N	%	95% CI	N	%	95% CI
Depressive episodes	2,265	71.2	62.9-79.5	492	15.5	9.1-21.8	188	5.9	2.1-9.7	83	2.6	0.0-5.3	44	1.4	0.0-2.7	110	3.5	0.3-6.6
Dysthymia	371	70.5	52.9-88.0	54	10.3	1.3-19.4	61	11.5	0.0-26.1	12	2.2	0.0-6.3	19	3.5	0.0-8.3	10	2.0	0.0-5.7
Bipolar affective disorder	1,204	85.9	73.8-98.0	108	7.7	0.0-16.2	10	0.7	0.0-2.2	25	1.8	0.0-4.5	20	1.4	0.0-3.5	35	2.5	0.0-5.6
Any affective disorder	3,631	76.3	68.9-83.8	588	12.4	7.3-17.4	210	4.4	1.7-7.2	120	2.5	0.4-4.6	63	1.3	0.3-2.4	145	3.0	0.7-5.4

Table B.11: Days partially unable to carry out their work, study or day-to-day activities in the previous four weeks, by type of 12-month affective disorder, subgrouped by weeks

ICD-10 affective disorder	0 days			1-7 days			8-14 days			15-21 days			22-28 days			Missing		
	N	%	95% CI	N	%	95% CI	N	%	95% CI	N	%	95% CI	N	%	95% CI	N	%	95% CI
Depressive episodes	1,661	52.2	41.9-62.5	898	28.2	19.5-37.0	296	9.3	4.6-14.0	126	4.0	0.6-7.3	103	3.2	1.1-5.4	98	3.1	0.0-6.1
Dysthymia	267	50.7	29.7-71.7	159	30.3	11.1-49.4	64	12.2	2.3-22.2	17	3.3	0.0-9.4	10	2.0	0.0-5.7	8	1.6	0.0-4.5
Bipolar affective disorder	990	70.7	48.6-92.7	209	14.9	1.7-28.2	116	8.3	0.0-17.0	20	1.4	0.0-3.5	31	2.2	0.0-5.1	35	2.5	0.0-5.6
Any affective disorder	2,800	58.9	48.0-69.7	1,111	23.3	15.6-31.1	433	9.1	4.9-13.3	146	3.1	0.7-5.4	134	2.8	1.1-4.5	132	2.8	0.5-5.0

Table B. 12: Four-week service use (times seen a doctor), by type of 12-month affective disorder (from K10 plus in relation to general symptoms of psychological distress reported in the K10)

ICD-10 affective disorder	0 times			1 time			2 times			3 times			4 times			5+ times			Missing		
	N	%	95% CI	N	%	95% CI	N	%	95% CI	N	%	95% CI	N	%	95% CI	N	%	95% CI	N	%	95% CI
Depressive episodes	1,947	61.2	51.6–70.7	289	9.1	3.3–14.8	283	8.9	4.2–13.6	131	4.1	0.8–7.4	231	7.3	3.2–11.3	191	6.0	2.0–10.0	111	3.5	0.4–6.6
Dysthymia	289	54.9	33.9–75.9	86	16.4	0.0–35.5	107	20.4	4.8–35.9	22	4.1	0.0–9.9	11	2.0	0.0–5.8	12	2.2	0.0–6.3	0	0.0	–
Bipolar affective disorder	1,130	80.6	64.8–96.4	86	6.1	0.0–13.5	38	2.7	0.0–5.9	0	0.0	–	73	5.2	0.0–12.1	40	2.8	0.0–6.2	35	2.5	0.0–5.6
Any affective disorder	3,146	66.1	56.5–75.8	431	9.1	4.0–14.1	346	7.3	3.7–10.8	152	3.2	0.8–5.6	293	6.2	2.7–9.6	242	5.1	2.1–8.1	146	3.1	0.7–5.4

Table B. 13: Days totally unable to carry out their work, study or day-to-day activities in the previous four weeks due to psychological distress, by type of 12-month anxiety disorder, subgrouped by weeks

ICD-10 anxiety disorder	0 days			1–7 days			8–14 days			15–21 days			22–28 days			Missing		
	N	%	95% CI	N	%	95% CI	N	%	95% CI	N	%	95% CI	N	%	95% CI	N	%	95% CI
Panic attack	2,419	68.4	58.3–78.4	747	21.1	11.3–30.9	171	4.8	2.0–7.6	104	2.9	1.1–4.8	31	0.9	0.0–1.9	66	1.9	0.1–3.6
Panic disorder	441	62.2	46.9–77.5	210	29.7	14.6–44.7	0	0.0	–	36	5.1	0.3–10.0	0	0.0	–	22	3.0	0.0–7.1
Agoraphobia	838	66.4	53.4–79.4	217	17.2	6.6–27.9	54	4.3	0.0–9.2	48	3.8	0.2–7.5	12	0.9	0.0–2.6	92	7.3	0.0–15.8
Social phobia	1,537	80.1	71.7–88.5	297	15.5	7.5–23.5	50	2.6	0.1–5.1	25	1.3	0.0–3.1	0	0.0	–	10	0.5	0.0–1.6
Specific phobia	2,279	75.7	66.9–84.5	337	11.2	6.0–16.4	232	7.7	0.6–14.8	98	3.3	0.2–6.3	23	0.8	0.0–1.8	41	1.4	0.0–3.4
Generalised anxiety disorder	331	62.1	43.5–80.7	141	26.5	8.2–44.8	29	5.4	0.0–11.4	0	0.0	–	12	2.2	0.0–6.2	20	3.8	0.0–8.9

ICD-10 anxiety disorder	0 days			1-7 days			8-14 days			15-21 days			22-28 days			Missing		
	N	%	95% CI	N	%	95% CI	N	%	95% CI	N	%	95% CI	N	%	95% CI	N	%	95% CI
Obsessive-compulsive disorder	1,315	83.1	67.6-98.7	90	5.7	0.0-11.8	143	9.1	0.0-21.9	22	1.4	0.0-3.4	12	0.7	0.0-2.2	0	0.0	-
Post-traumatic stress disorder	3,220	77.2	68.2-86.3	508	12.2	7.0-17.3	64	1.5	0.3-2.8	106	2.6	0.3-4.8	30	0.7	0.0-1.5	241	5.8	0.0-12.6
Any anxiety disorder	5,775	77.8	71.2-84.5	880	11.9	7.9-15.8	233	3.1	0.3-6.0	147	2.0	0.6-3.4	42	0.6	0.0-1.1	343	4.6	0.4-8.8

Table B. 14: Days partially out of role in the previous four weeks due to psychological distress, by type of 12-month anxiety disorder, subgrouped by weeks

ICD-10 anxiety disorder	0 days			1-7 days			8-14 days			15-21 days			22-28 days			Missing		
	N	%	95% CI	N	%	95% CI	N	%	95% CI	N	%	95% CI	N	%	95% CI	N	%	95% CI
Panic attack	1,869	52.8	41.6-64.0	1,026	29.0	18.6-39.5	299	8.4	4.4-12.5	195	5.5	2.6-8.4	103	2.9	1.0-4.9	45	1.3	0.0-2.8
Panic disorder	340	48.0	32.2-63.7	210	29.7	14.6-44.7	79	11.2	3.7-18.7	35	4.9	0.2-9.6	33	4.7	0.0-9.8	12	1.6	0.0-4.7
Agoraphobia	586	46.5	32.5-60.5	410	32.5	19.6-45.4	157	12.5	5.3-19.7	14	1.1	0.0-3.1	12	0.9	0.0-2.6	82	6.5	0.0-14.9
Social phobia	1,002	52.2	41.2-63.3	628	32.7	22.0-43.5	153	8.0	3.4-12.6	99	5.2	0.2-10.1	37	2.0	0.0-4.2	0	0.0	-
Specific phobia	1,820	60.4	50.3-70.6	811	26.9	17.4-36.5	169	5.6	2.5-8.7	88	2.9	0.0-5.9	51	1.7	0.2-3.2	72	2.4	0.0-5.2
Generalised anxiety disorder	235	44.0	25.2-62.9	124	23.2	6.2-40.2	121	22.7	5.9-39.6	15	2.7	0.0-6.4	39	7.3	0.0-15.4	0	0.0	-
Obsessive-compulsive disorder	1,126	71.2	50.0-92.5	318	20.1	2.9-37.4	65	4.1	0.1-8.1	20	1.3	0.0-3.2	51	3.2	0.0-8.6	0	0.0	-
Post-traumatic stress disorder	2,228	53.4	40.2-66.7	1,368	32.8	21.7-43.9	262	6.3	2.6-10.0	175	4.2	1.3-7.1	54	1.3	0.2-2.4	82	2.0	0.0-4.1
Any anxiety disorder	4,351	58.6	49.5-67.8	2,092	28.2	20.7-35.7	424	5.7	3.2-8.2	234	3.2	1.4-4.9	155	2.1	0.6-3.5	164	2.2	0.2-4.2

Table B. 15: Estimated 12-month service use, by type of 12-month anxiety disorder (from CID) in relation to the type of affective disorder reported)

ICD-10 disorder	Received professional treatment in last 12 months																			
	Yes					No					Don't know					Refused				
	N	%	95% CI	N	%	95% CI	N	%	95% CI	N	%	95% CI	N	%	95% CI	N	%	95% CI		
Panic disorder	343	48.3	32.7–63.9	367	51.7	36.1–67.3	0	0.0	–	0	0.0	–	0	0.0	–	0	0.0	–		
Agoraphobia	584	46.3	32.7–60.0	677	53.7	40.0–67.3	0	0.0	–	0	0.0	–	0	0.0	–	0	0.0	–		
Social phobia	484	25.2	16.1–34.3	1,435	74.8	65.7–83.9	0	0.0	–	0	0.0	–	0	0.0	–	0	0.0	–		
Specific phobia	371	12.3	6.0–18.6	2,640	87.7	81.4–94.0	0	0.0	–	0	0.0	–	0	0.0	–	0	0.0	–		
Generalised anxiety disorder	403	75.6	60.0–91.2	130	24.4	8.8–40.0	0	0.0	–	0	0.0	–	0	0.0	–	0	0.0	–		
Obsessive-compulsive disorder	185	11.7	2.5–20.9	1,349	85.3	73.6–97.0	0	0.0	–	0	0.0	–	47	3.0	0.0–8.8					
Post-traumatic stress disorder ^a	1,068	50.2	38.7–61.7	1,041	48.9	37.4–60.4	7	0.3	0.0–0.9	12	0.5	0.0–1.6								

^a Does not cover all members with this disorder, but percentages are out of the total number with responses.

Table B. 16: Four-week service use (times seen a doctor), by type of 12-month anxiety disorder (from K10 in relation to general symptoms of psychological distress reported in the K10)

ICD-10 anxiety disorder	0 times		1 time		2 times		3 times		4 times		5+ times		Missing	
	N	95% CI	N	95% CI	N	95% CI	N	95% CI	N	95% CI	N	95% CI	N	95% CI
Panic attack	2,036	57.6 46.5–68.6	442	12.5 2.6–22.5	434	12.3 6.4–18.2	169	4.8 1.2–8.3	161	4.6 2.0–7.2	214	6.1 2.9–9.2	80	2.3 0.4–4.2
Panic disorder	358	50.5 34.8–66.2	48	6.7 1.0–12.5	152	21.5 6.7–36.3	14	1.9 0.0–5.5	46	6.5 0.0–14.1	45	6.3 0.9–11.6	47	6.6 0.4–12.8
Agoraphobia	647	51.3 37.5–65.1	83	6.6 0.0–13.3	249	19.7 8.0–31.5	0	0.0 –	75	5.9 0.6–11.2	114	9.0 2.1–16.0	93	7.4 0.0–15.9
Social phobia	1,309	68.2 58.0–78.4	85	4.4 0.8–8.1	293	15.3 7.0–23.6	68	3.6 0.0–8.3	50	2.6 0.1–5.1	113	5.9 1.3–10.5	0	0.0 –
Specific phobia	2,165	71.9 62.7–81.1	110	3.6 0.7–6.6	184	6.1 2.1–10.1	138	4.6 0.3–8.9	214	7.1 0.2–14.1	143	4.7 1.7–7.8	57	1.9 0.0–4.2

ICD-10 anxiety disorder	0 times		1 time		2 times		3 times		4 times		5+ times		Missing								
	N	%	95% CI	N	%	95% CI	N	%	95% CI	N	%	95% CI	N	%	95% CI						
Generalised anxiety disorder	256	48.0	29.0–67.1	71	13.2	1.3–25.2	118	22.0	5.3–38.8	0	0.0	–	20	3.8	0.0–9.1	58	10.9	0.0–23.2	10	1.9	0.0–5.4
Obsessive-compulsive disorder	1,145	72.4	51.6–93.2	61	3.9	0.0–9.0	103	6.5	0.7–12.3	101	6.4	0.0–14.7	148	9.4	0.0–22.5	22	1.4	0.0–3.4	0	0.0	–
Post-traumatic stress disorder	2,867	68.8	58.1–79.5	254	6.1	3.0–9.2	362	8.7	3.9–13.5	151	3.6	0.9–6.3	131	3.1	0.7–5.6	150	3.6	0.9–6.3	255	6.1	0.0–13.0
Any anxiety disorder	5,085	68.5	60.5–76.5	390	5.3	3.0–7.5	731	9.9	6.1–13.6	246	3.3	1.1–5.5	331	4.5	1.3–7.6	276	3.7	1.7–5.7	360	4.9	0.6–9.1

Table B.17: Total days out of role in the previous four weeks due to psychological distress, by type of 12-month alcohol disorder, subgrouped by weeks

ICD-10 alcohol disorder	0 days		1–7 days		8–14 days		15–21 days		22–28 days		Missing							
	N	%	95% CI	N	%	95% CI	N	%	95% CI	N	%	95% CI						
Alcohol harmful use	1,266	89.2	80.8–97.5	61	4.3	0.0–9.2	23	1.6	0.0–3.9	20	1.4	0.0–3.5	0	0.0	–	49	3.5	0.0–8.9
Alcohol dependence	975	83.3	73.1–93.5	113	9.6	1.8–17.5	19	1.6	0.0–3.7	0	0.0	–	0	0.0	–	64	5.5	0.0–12.4
Any alcohol disorder	2,241	86.5	79.8–93.2	174	6.7	2.2–11.3	42	1.6	0.0–3.2	20	0.8	0.0–1.9	0	0.0	–	113	4.4	0.0–8.7

Table B.18: Partial days out of role in the previous four weeks due to psychological distress, by type of 12-month alcohol disorder, subgrouped by weeks

ICD-10 alcohol disorder	0 days		1–7 days		8–14 days		15–21 days		22–28 days		Missing							
	N	%	95% CI	N	%	95% CI	N	%	95% CI	N	%	95% CI						
Alcohol harmful alcohol use	1,176	82.9	72.5–93.2	151	10.7	3.3–18.0	32	2.3	0.0–4.9	0	0.0	–	20	1.4	0.0–3.3	40	2.8	0.0–8.1
Alcohol dependence	774	66.1	52.6–79.5	156	13.4	4.8–21.9	133	11.3	1.3–21.4	25	2.2	0.0–4.6	10	0.9	0.0–2.5	72	6.2	0.0–13.2
Any alcohol disorder	1,950	75.3	66.2–84.4	308	11.9	6.2–17.5	165	6.4	1.3–11.4	25	1.0	0.0–2.1	30	1.2	0.0–2.5	112	4.3	0.0–8.7

Table B.19: Four-week service use (number of visits to the doctor), by type of 12-month alcohol disorder (from K10 in relation to general symptoms of psychological distress reported in the K10)

ICD-10 alcohol disorder	0 times			1 time			2 times			3 times			4 times			5+ times			Missing		
	N	%	95% CI	N	%	95% CI	N	%	95% CI	N	%	95% CI	N	%	95% CI	N	%	95% CI	N	%	95% CI
Alcohol harmful use	1,183	83.4	72.5–94.2	19	1.3	0.0–3.2	91	6.4	0.0–13.6	12	0.8	0.0–2.4	32	2.3	0.0–4.9	20	1.4	0.0–3.5	63	4.4	0.0–10.2
Alcohol dependence	808	69.0	55.5–82.5	111	9.5	1.6–17.3	19	1.6	0.0–3.9	29	2.5	0.0–5.9	41	3.5	0.0–10.0	90	7.7	0.0–15.8	72	6.2	0.0–13.2
Any alcohol disorder	1,991	76.9	67.8–86.0	130	5.0	1.2–8.9	110	4.2	0.2–8.2	41	1.6	0.0–3.4	73	2.8	0.0–6.1	110	4.3	0.3–8.3	135	5.2	0.7–9.7

Table B.20: Estimated prevalence of ICD-10 post-traumatic stress disorder among those reporting different traumatic event types in the CIDI

Traumatic event	Total no. experiencing the event			ICD-10 PTSD associated with each event		
	N	%	95% CI	N	%	95% CI
Saw someone badly injured/killed or unexpectedly saw a dead body	22,204	44.4	40.5–48.3	3,057	13.8	9.3–18.3
Someone close died	18,159	36.3	32.7–39.8	2,407	13.3	8.2–18.4
Peacekeeper (in war zone or place of ongoing terror)	15,781	31.5	27.9–35.2	1,456	9.2	6.3–12.1
Combat (military or organised non-military group)	14,941	29.9	26.4–33.3	1,550	10.4	7.5–13.3
Life-threatening automobile accident	13,395	26.8	23.1–30.5	1,667	12.4	5.5–19.4
Mugged, held up, threatened with weapon	12,994	26.0	22.4–29.5	2,303	17.7	10.3–25.1
Exposed to toxic chemical that could cause serious harm	11,390	22.8	19.1–26.4	1,527	13.4	5.3–21.5
Major natural disaster	8,862	17.7	15.3–20.2	1,148	13.0	9.2–16.7
Other life-threatening accident	7,815	15.6	12.3–18.9	1,425	18.2	6.7–29.7
Man-made disaster	7,376	14.7	11.9–17.6	1,371	18.6	7.4–29.8
Saw atrocities or carnage such as mutilated bodies or mass killings	6,923	13.8	11.1–16.6	1,541	22.3	10.7–33.9
Badly beaten by anyone else	6,745	13.5	10.6–16.3	1,106	16.4	4.0–28.8
Witnessed serious physical fights at home as a child	6,690	13.4	10.5–16.2	1,268	19.0	6.3–31.6
Life-threatening illness	6,137	12.3	9.2–15.3	588	9.6	5.6–13.6
Other traumatic event	6,034	12.1	9.4–14.7	1,638	27.1	14.1–40.1
Someone close had traumatic experience	5,677	11.3	9.2–13.4	754	13.3	9.1–17.5
Sexual assault	4,792	9.6	7.2–12.0	548	11.4	7.1–15.8
Child had life-threatening illness/injury	3,172	6.3	4.8–7.8	526	16.6	9.6–23.6
Stalked	2,659	5.3	3.2–7.4	1,020	38.4	15.1–61.7
Raped	2,518	5.0	2.9–7.1	1,066	42.3	18.5–66.1
Accidentally injured/killed someone	2,293	4.6	1.9–7.3	651	28.4	0.0–62.0
Beaten by parents/guardian as a child	2,291	4.6	3.4–5.7	401	17.5	10.0–25.1
Experience don't want to talk about	2,131	4.3	3.3–5.2	574	26.9	17.8–36.1
Unarmed civilian in a place of war, revolution, military coup or invasion	1,886	3.8	2.6–4.9	344	18.3	4.1–32.4
Purposefully injured/tortured/killed someone	1,614	3.2	1.8–4.6	267	16.5	5.8–27.3
Lived as a civilian in a place of ongoing terror for political, ethnic, religious or other reasons	1,305	2.6	1.7–3.5	46	3.5	0.3–6.7
Kidnapped or held captive	864	1.7	0.0–3.7	678	78.5	52.3–100.0
Beaten by spouse/romantic partner	770	1.5	1.0–2.1	240	31.1	14.2–48.1
Refugee	74	0.1	0.0–0.3	8	11.0	0.0–34.3

Section 2 data tables

Table B.21: K10 risk categories (two categories) by sex, Service, sex by Service, rank and deployment status

	N	Low (10–19)			High (20–50)		
		N	%	% CI	N	%	% CI
Total	50,049	41,008	81.9	81.5–82.4	9,041	18.1	17.6–18.5
Males	43,241	35,658	82.5	82.0–83.0	7,583	17.5	17.0–18.0
Navy	9,508	7,562	79.5	78.4–80.7	1,946	20.5	19.3–21.6
Army	22,843	18,852	82.5	81.8–83.3	3,991	17.5	16.7–18.2
Air Force	10,890	9,245	84.9	84.2–85.6	1,645	15.1	14.4–15.8
Females	6,808	5,351	78.6	77.5–79.7	1,457	21.4	20.3–22.5
Navy	2,104	1,559	74.1	71.7–76.5	545	25.9	23.5–28.3
Army	2,513	2,002	79.7	77.9–81.4	511	20.3	18.6–22.1
Air Force	2,191	1,790	81.7	80.1–83.2	401	18.3	16.8–19.9
Navy	11,612	9,120	78.5	77.5–79.6	2,492	21.5	20.4–22.5
Army	25,356	20,854	82.2	81.6–82.9	4,502	17.8	17.1–18.4
Air Force	13,081	11,034	84.4	83.7–85.0	2,047	15.6	15.0–16.3
Officers	12,034	10,311	85.7	85.1–86.3	1,723	14.3	13.7–14.9
Non-commissioned officers	22,319	18,472	82.8	82.2–83.3	3,847	17.2	16.7–17.8
Other ranks	15,696	12,226	77.9	76.7–79.0	3,470	22.1	21.0–23.3
Never deployed	20,171	15,866	78.7	77.8–79.5	4,305	21.3	20.5–22.2
Deployed at least once	29,878	25,143	84.2	83.6–84.7	4,735	15.8	15.3–16.4

Table B.22: K10 risk categories, by sex by Service and by Service, rank and deployment status

	N	Low (10–15)			Moderate (16–21)			High (22–29)			Very high (30–50)		
		N	%	% CI	N	%	% CI	N	%	% CI	N	%	% CI
Total	50,049	32,380	64.7	64.1–65.3	11,237	22.5	22.0–22.9	4,655	9.3	9.0–9.6	1,778	3.6	3.3–3.8
Males	43,241	28,397	65.7	65.1–66.3	9,504	22.0	21.4–22.5	3,909	9.0	8.7–9.4	1,431	3.3	3.1–3.6
Navy	9,508	5,810	61.1	59.7–62.5	2,347	24.7	23.4–25.9	979	10.3	9.4–11.2	372	3.9	3.3–4.5
Army	22,843	15,178	66.4	65.5–67.4	4,788	21.0	20.2–21.8	2,093	9.2	8.6–9.7	784	3.4	3.1–3.8
Air Force	10,890	7,409	68.0	67.1–68.9	2,369	21.8	20.9–22.6	836	7.7	7.2–8.2	275	2.5	2.2–2.8
Females	6,808	3,983	58.5	57.2–59.8	1,733	25.5	24.3–26.6	746	11.0	10.1–11.8	347	5.1	4.5–5.7
Navy	2,104	1,156	54.9	52.2–57.6	537	25.5	23.2–27.9	268	12.7	10.9–14.6	144	6.8	5.4–8.2
Army	2,513	1,505	59.9	57.8–62.0	625	24.9	23.0–26.7	266	10.6	9.2–11.9	118	4.7	3.7–5.6
Air Force	2,191	1,322	60.3	58.4–62.3	571	26.1	24.3–27.8	212	9.7	8.5–10.9	86	3.9	3.1–4.7
Navy	11,612	6,966	60.0	58.7–61.2	2,884	24.8	23.7–25.9	1,247	10.7	9.9–11.5	516	4.4	3.9–5.0
Army	25,356	16,683	65.8	65.0–66.6	5,413	21.3	20.6–22.1	2,359	9.3	8.8–9.8	901	3.6	3.2–3.9
Air Force	13,081	8,731	66.7	65.9–67.6	2,940	22.5	21.7–23.2	1,049	8.0	7.5–8.5	361	2.8	2.5–3.0
Officers	12,034	8,353	69.4	68.6–70.2	2,500	20.8	20.1–21.5	930	7.7	7.3–8.2	251	2.1	1.9–2.3
Non-commissioned officers	22,319	14,723	66.0	65.3–66.6	4,878	21.9	21.3–22.5	1,977	8.9	8.5–9.3	741	3.3	3.1–3.6
Other ranks	15,696	9,304	59.3	57.9–60.6	3,859	24.6	23.4–25.8	1,747	11.1	10.3–12.0	786	5.0	4.4–5.6
Never deployed	20,170	11,909	59.0	58.1–60.0	5,228	25.9	25.0–26.8	2,174	10.8	10.2–11.4	859	4.3	3.8–4.7
Deployed at least once	29,878	20,471	68.5	67.8–69.2	6,008	20.1	19.5–20.7	2,481	8.3	7.9–8.7	918	3.1	2.8–3.3

Table B.23: Days in the previous four weeks ADF personnel were totally unable to carry out their work, study or day-to-day activities due to psychological distress (subgrouped by weeks)

	N	Missing ^a			0 days			1-7 days			8-14 days			15-21 days			22-28 days		
		N	%	% CI	N	%	% CI	N	%	% CI	N	%	% CI	N	%	% CI	N	%	% CI
Total	50,049	1,885	3.8	3.5-4.0	44,884	89.7	89.3-90.0	2,644	5.3	5.0-5.5	309	0.6	0.5-0.7	178	0.4	0.3-0.4	148	0.3	0.2-0.4
Males	43,241	1,647	3.8	3.5-4.1	39,039	90.3	89.9-90.7	2,014	4.7	4.4-4.9	258	0.6	0.5-0.7	154	0.4	0.3-0.4	129	0.3	0.2-0.4
Navy	9,508	256	2.7	2.2-3.1	8,649	91.0	90.1-91.8	494	5.2	4.5-5.8	41	0.4	0.2-0.6	40	0.4	0.2-0.6	29	0.3	0.2-0.4
Army	22,843	1,081	4.7	4.3-5.2	20,358	89.1	88.5-89.7	1,079	4.7	4.3-5.1	148	0.6	0.5-0.8	98	0.4	0.3-0.6	78	0.3	0.2-0.5
Air Force	10,890	310	2.8	2.5-3.2	10,032	92.1	91.6-92.7	441	4.1	3.7-4.4	69	0.6	0.5-0.8	17	0.2	0.1-0.2	21	0.2	0.1-0.3
Females	6,808	238	3.5	3.0-4.0	5,845	85.9	84.9-86.8	630	9.3	8.5-10.0	52	0.8	0.5-1.0	23	0.3	0.2-0.5	20	0.3	0.1-0.4
Navy	2,104	78	3.7	2.6-4.8	1,764	83.9	81.9-85.9	223	10.6	8.9-12.3	11	0.5	0.2-0.9	13	0.6	0.2-1.1	15	0.7	0.3-1.1
Army	2,513	78	3.1	2.3-3.9	2,167	86.2	84.7-87.7	234	9.3	8.0-10.6	27	1.1	0.6-1.5	6	0.2	0.0-0.4	2	0.1	0.0-0.2
Air Force	2,191	83	3.8	3.0-4.6	1,914	87.4	86.0-88.7	173	7.9	6.8-9.0	14	0.6	0.3-1.0	4	0.2	0.0-0.4	3	0.1	0.0-0.3
Navy	11,612	333	2.9	2.5-3.3	10,413	89.7	88.9-90.4	716	6.2	5.6-6.8	52	0.5	0.3-0.6	53	0.5	0.3-0.6	44	0.4	0.2-0.5
Army	25,356	1,159	4.6	4.2-5.0	22,525	88.8	88.3-89.4	1,313	5.2	4.8-5.6	174	0.7	0.5-0.8	105	0.4	0.3-0.5	80	0.3	0.2-0.4
Air Force	13,081	393	3.0	2.7-3.3	11,946	91.3	90.8-91.8	615	4.7	4.3-5.1	83	0.6	0.5-0.8	21	0.2	0.1-0.2	24	0.2	0.1-0.3
Officers	12,034	251	2.1	1.8-2.3	11,200	93.1	92.6-93.5	497	4.1	3.8-4.5	47	0.4	0.3-0.5	18	0.1	0.1-0.2	21	0.2	0.1-0.2
Non-commissioned officers	22,319	848	3.8	3.5-4.1	19,995	89.6	89.1-90.0	1,200	5.4	5.1-5.7	140	0.6	0.5-0.7	75	0.3	0.3-0.4	61	0.3	0.2-0.3
Other ranks	15,696	787	5.0	4.4-5.7	13,690	87.2	86.3-88.1	948	6.0	5.4-6.7	122	0.8	0.5-1.0	85	0.5	0.3-0.7	66	0.4	0.3-0.6
Never deployed	20,171	596	3.0	2.6-3.3	18,071	89.6	89.0-90.2	1,220	6.0	5.6-6.5	135	0.7	0.5-0.8	89	0.4	0.3-0.6	59	0.3	0.2-0.4
Deployed at least once	29,878	1,289	4.3	4.0-4.6	26,813	89.7	89.3-90.2	1,424	4.8	4.5-5.1	175	0.6	0.5-0.7	89	0.3	0.2-0.4	89	0.3	0.2-0.4

^a Responded to all K10 questions but did not respond to this question.

Table B.24: Days in the previous four weeks ADF personnel were partially unable to carry out their work, study or day-to-day activities due to psychological distress (subgrouped by weeks)

	N	Missing ^a			0 days			1-7 days			8-14 days			15-21 days			22-28 days		
		N	%	% CI	N	%	% CI	N	%	% CI	N	%	% CI	N	%	% CI	N	%	% CI
Total	50,049	2,034	4.1	3.8-4.3	40,478	80.9	80.4-81.3	5,290	10.6	10.2-10.9	1,056	2.1	2.0-2.3	589	1.2	1.1-1.3	603	1.2	1.1-1.3
Males	43,241	1,764	4.1	3.8-4.4	35,394	81.9	81.4-82.3	4,286	9.9	9.5-10.3	825	1.9	1.7-2.1	481	1.1	1.0-1.2	491	1.1	1.0-1.3
Navy	9,508	272	2.9	2.4-3.3	7,722	81.2	80.1-82.3	1,068	11.2	10.3-12.1	204	2.1	1.7-2.6	113	1.2	0.9-1.5	130	1.4	1.0-1.7
Army	22,843	1,156	5.1	4.6-5.5	18,705	81.9	81.1-82.6	2,067	9.0	8.5-9.6	410	1.8	1.6-2.0	250	1.1	0.9-1.3	255	1.1	0.9-1.3
Air Force	10,890	336	3.1	2.7-3.4	8,968	82.4	81.6-83.1	1,151	10.6	10.0-11.2	212	1.9	1.7-2.2	118	1.1	0.9-1.3	106	1.0	0.8-1.2
Females	6,808	270	4.0	3.4-4.5	5,084	74.7	73.5-75.8	1,004	14.7	13.8-15.7	231	3.4	2.9-3.9	107	1.6	1.2-1.9	112	1.7	1.3-2.0
Navy	2,104	89	4.2	3.1-5.4	1,514	72.0	69.6-74.4	353	16.8	14.8-18.8	84	4.0	2.9-5.1	29	1.4	0.7-2.0	35	1.7	1.0-2.3
Army	2,513	99	3.9	3.1-4.8	1,902	75.7	73.8-77.5	326	13.0	11.6-14.4	81	3.2	2.4-4.0	48	1.9	1.3-2.5	57	2.3	1.6-3.0
Air Force	2,191	82	3.7	3.0-4.5	1,688	76.1	74.4-77.8	325	14.8	13.4-16.2	66	3.0	2.3-3.7	31	1.4	0.9-1.9	20	0.9	0.5-1.3
Navy	11,612	361	3.1	2.7-3.6	9,236	79.5	78.5-80.5	1,421	12.2	11.4-13.0	288	2.5	2.1-2.9	142	1.2	0.9-1.5	164	1.4	1.1-1.7
Army	25,356	1,255	4.9	4.5-5.4	20,606	81.3	80.6-82.0	2,393	9.4	8.9-9.9	491	1.9	1.7-2.2	298	1.2	1.0-1.4	313	1.2	1.0-1.4
Air Force	13,081	418	3.2	2.9-3.5	10,636	81.3	80.6-82.0	1,475	11.3	10.7-11.8	278	2.1	1.9-2.4	149	1.1	1.0-1.3	126	1.0	0.8-1.1
Officers	12,034	274	2.3	2.0-2.5	9,880	82.1	81.5-82.7	1,486	12.3	11.8-12.9	197	1.6	1.4-1.8	106	0.9	0.7-1.0	92	0.8	0.6-0.9
Non-commissioned officers	22,319	908	4.1	3.8-4.4	18,027	80.8	80.2-81.3	2,270	10.2	9.7-10.6	526	2.4	2.1-2.6	274	1.2	1.1-1.4	314	1.4	1.2-1.6
Other ranks	15,696	853	5.4	4.8-6.1	12,570	80.1	79.0-81.2	1,533	9.8	9.0-10.6	334	2.1	1.8-2.5	209	1.3	1.0-1.6	197	1.3	1.0-1.6
Never deployed	20,170	642	3.2	2.8-3.6	16,267	80.6	79.9-81.4	2,323	11.5	10.9-12.1	454	2.3	2.0-2.5	245	1.2	1.0-1.4	239	1.2	1.0-1.4
Deployed at least once	29,879	1,392	4.7	4.3-5.0	24,211	81.0	80.5-81.6	2,966	9.9	9.5-10.3	602	2.0	1.8-2.2	344	1.2	1.0-1.3	364	1.2	1.1-1.4

^a Responded to all K10 questions but did not respond to this question.

Table B.25: Number of times ADF personnel reported seeing a doctor in the previous four weeks for symptoms of psychological distress reported in the K10

	N	Missing ^a		Did not seek help		Sought help once		Sought help twice		Sought help three times		Sought help four times		Sought help more than four times	
		N	%	N	%	N	%	N	%	N	%	N	%	N	%
Total	50,049	1,942	3.9	43,870	87.7	1,708	3.4	1,051	2.1	506	1.0	396	0.8	576	1.2
Males	43,241	1,686	3.9	38,259	88.5	1,347	3.1	790	1.8	417	1.0	315	0.7	427	1.0
Navy	9,508	282	3.0	8,487	89.3	323	3.4	195	2.0	91	1.0	55	0.6	75	0.8
Army	22,843	1,072	4.7	19,985	87.5	636	2.8	436	1.9	237	1.0	195	0.9	282	1.2
Air Force	10,890	331	3.0	9,787	89.9	389	3.6	160	1.5	89	0.8	65	0.6	69	0.6
Females	6,808	257	3.8	5,611	82.4	360	5.3	261	3.8	89	1.3	81	1.2	149	2.2
Navy	2,104	87	4.1	1,725	82.0	97	4.6	84	4.0	29	1.4	37	1.7	46	2.2
Army	2,513	89	3.6	2,067	82.3	137	5.4	92	3.7	39	1.5	25	1.0	64	2.5
Air Force	2,191	80	3.7	1,818	83.0	127	5.8	85	3.9	22	1.0	19	0.9	40	1.8
Navy	11,612	369	3.2	10,212	87.9	420	3.6	279	2.4	120	1.0	92	0.8	121	1.0
Army	25,356	1,162	4.6	22,053	87.0	772	3.0	527	2.1	275	1.1	220	0.9	346	1.4
Air Force	13,081	411	3.1	11,606	88.7	516	3.9	244	1.9	111	0.8	84	0.6	109	0.8
Officers	12,034	267	2.2	10,969	91.2	365	3.0	187	1.6	123	1.0	48	0.4	74	0.6
Non-commissioned officers	22,319	859	3.9	19,497	87.4	806	3.6	499	2.2	194	0.9	192	0.9	271	1.2
Other ranks	15,696	816	5.2	13,404	85.4	537	3.4	365	2.3	189	1.2	156	1.0	231	1.5
Never deployed	20,170	638	3.2	17,779	88.1	670	3.3	491	2.4	222	1.1	132	0.7	238	1.2
Ever deployed	29,879	1,304	4.4	26,091	87.3	1,038	3.5	559	1.9	284	1.0	264	0.9	338	1.1

^a Responded to all K10 questions but did not respond to this question.

Table B.26: Properties of the K10 for predicting 30-day ICD-10 anxiety disorder

Cut-off	Sensitivity		Specificity		Positive predictive value		Negative predictive value	
	Value	95% CI	Value	95% CI	Value	95% CI	Value	95% CI
10	1.00	1.00–1.00	0.00	0.00–0.00	0.08	0.05–0.10	–	–
11	1.00	0.99–1.00	0.14	0.11–0.18	0.09	0.06–0.11	1.00	0.99–1.00
12	1.00	0.99–1.00	0.27	0.22–0.31	0.10	0.07–0.13	1.00	1.00–1.00
13	0.84	0.62–1.00	0.36	0.32–0.41	0.10	0.08–0.12	0.97	0.91–1.00
14	0.80	0.58–1.00	0.46	0.42–0.51	0.11	0.09–0.13	0.97	0.92–1.00
15	0.77	0.57–0.98	0.55	0.51–0.59	0.12	0.10–0.15	0.97	0.93–1.00
16	0.73	0.53–0.93	0.64	0.60–0.68	0.14	0.12–0.17	0.97	0.93–1.00
17	0.68	0.49–0.87	0.72	0.68–0.75	0.16	0.13–0.20	0.96	0.94–0.99
18	0.61	0.44–0.79	0.76	0.72–0.79	0.17	0.14–0.21	0.96	0.93–0.99
19	0.58	0.41–0.75	0.81	0.78–0.84	0.20	0.16–0.24	0.96	0.93–0.99
20	0.49	0.35–0.64	0.84	0.82–0.87	0.21	0.17–0.25	0.95	0.93–0.98
21	0.45	0.31–0.58	0.87	0.85–0.89	0.22	0.18–0.27	0.95	0.93–0.98
22	0.42	0.29–0.54	0.89	0.88–0.91	0.24	0.19–0.30	0.95	0.92–0.97
23	0.37	0.25–0.49	0.91	0.89–0.92	0.25	0.19–0.31	0.95	0.92–0.97
24	0.35	0.23–0.46	0.92	0.91–0.94	0.27	0.21–0.34	0.95	0.92–0.97
25	0.31	0.21–0.42	0.93	0.92–0.95	0.27	0.20–0.34	0.94	0.92–0.97
26	0.30	0.19–0.40	0.95	0.93–0.96	0.31	0.23–0.39	0.94	0.92–0.97
27	0.25	0.16–0.34	0.95	0.94–0.96	0.30	0.21–0.38	0.94	0.92–0.96
28	0.24	0.15–0.33	0.96	0.95–0.97	0.34	0.24–0.43	0.94	0.92–0.96
29	0.21	0.13–0.29	0.98	0.97–0.98	0.41	0.31–0.52	0.94	0.92–0.96
30	0.19	0.12–0.27	0.98	0.97–0.99	0.43	0.32–0.54	0.94	0.91–0.96
31	0.17	0.10–0.24	0.98	0.98–0.99	0.45	0.33–0.57	0.94	0.91–0.96
32	0.13	0.08–0.19	0.99	0.98–0.99	0.44	0.31–0.57	0.93	0.91–0.96
33	0.11	0.06–0.17	0.99	0.98–0.99	0.44	0.30–0.58	0.93	0.91–0.95
34	0.11	0.06–0.16	0.99	0.99–0.99	0.48	0.33–0.64	0.93	0.91–0.95
35	0.10	0.05–0.15	0.99	0.99–1.00	0.49	0.32–0.66	0.93	0.91–0.95
36	0.09	0.04–0.13	0.99	0.99–1.00	0.52	0.33–0.70	0.93	0.91–0.95
37	0.07	0.03–0.11	0.99	0.99–1.00	0.50	0.30–0.69	0.93	0.91–0.95
38	0.04	0.02–0.07	0.99	0.99–1.00	0.41	0.21–0.60	0.93	0.90–0.95
39	0.04	0.01–0.06	0.99	0.99–1.00	0.36	0.16–0.55	0.93	0.90–0.95
40	0.03	0.01–0.05	1.00	0.99–1.00	0.44	0.20–0.68	0.93	0.90–0.95
41	0.02	0.01–0.04	1.00	1.00–1.00	0.40	0.14–0.67	0.93	0.90–0.95
42	0.01	0.00–0.03	1.00	1.00–1.00	0.33	0.05–0.60	0.93	0.90–0.95
43	0.01	0.00–0.03	1.00	1.00–1.00	0.35	0.05–0.65	0.93	0.90–0.95
44	0.01	0.00–0.02	1.00	1.00–1.00	0.28	0.04–0.60	0.92	0.90–0.95

Cut-off	Sensitivity		Specificity		Positive predictive value		Negative predictive value	
	Value	95% CI	Value	95% CI	Value	95% CI	Value	95% CI
45	0.01	0.00–0.01	1.00	1.00–1.00	0.25	0.11–0.61	0.92	0.90–0.95
46	0.01	0.00–0.01	1.00	1.00–1.00	0.51	0.03–0.99	0.92	0.90–0.95
47	0.00	0.00–0.00	1.00	1.00–1.00	0.00	0.00–0.00	0.92	0.90–0.95
48	0.00	0.00–0.00	1.00	1.00–1.00	0.00	0.00–0.00	0.92	0.90–0.95
49	0.00	0.00–0.00	1.00	1.00–1.00	0.00	0.00–0.00	0.92	0.90–0.95
50	0.00	0.00–0.00	1.00	1.00–1.00	0.00	0.00–0.00	0.92	0.90–0.95

AUC	
Value	95% CI
0.75	0.60–0.89

Table B.27: Properties of the K10 for predicting 30-day ICD-10 affective disorder

Cut-off	Sensitivity		Specificity		Positive predictive value		Negative predictive value	
	Value	95% CI	Value	95% CI	Value	95% CI	Value	95% CI
10	1.00	1.00–1.00	0.00	0.00–0.00	0.03	0.02–0.03	–	–
11	1.00	1.00–1.00	0.14	0.11–0.17	0.03	0.02–0.04	1.00	1.00–1.00
12	1.00	1.00–1.00	0.25	0.21–0.29	0.04	0.03–0.05	1.00	1.00–1.00
13	0.88	0.73–1.00	0.36	0.31–0.40	0.04	0.03–0.05	0.99	0.98–1.00
14	0.88	0.73–1.03	0.45	0.41–0.50	0.04	0.03–0.05	0.99	0.98–1.00
15	0.80	0.64–0.96	0.54	0.50–0.58	0.04	0.03–0.06	0.99	0.98–1.00
16	0.80	0.64–0.96	0.62	0.59–0.66	0.05	0.04–0.07	0.99	0.98–1.00
17	0.78	0.62–0.94	0.70	0.66–0.73	0.07	0.05–0.08	0.99	0.98–1.00
18	0.76	0.60–0.92	0.74	0.71–0.77	0.07	0.05–0.09	0.99	0.98–1.00
19	0.75	0.59–0.91	0.79	0.76–0.82	0.09	0.06–0.11	0.99	0.98–1.00
20	0.69	0.54–0.85	0.83	0.81–0.85	0.10	0.07–0.13	0.99	0.98–1.00
21	0.62	0.46–0.77	0.86	0.84–0.88	0.11	0.08–0.14	0.99	0.98–1.00
22	0.58	0.43–0.73	0.88	0.87–0.90	0.12	0.08–0.15	0.99	0.98–0.99
23	0.54	0.40–0.69	0.90	0.88–0.91	0.13	0.09–0.17	0.99	0.98–0.99
24	0.54	0.39–0.69	0.91	0.90–0.93	0.15	0.10–0.19	0.99	0.98–0.99
25	0.44	0.31–0.58	0.92	0.91–0.94	0.14	0.09–0.18	0.98	0.98–0.99
26	0.41	0.27–0.54	0.94	0.92–0.95	0.15	0.10–0.20	0.98	0.98–0.99
27	0.37	0.24–0.49	0.94	0.93–0.96	0.15	0.09–0.21	0.98	0.98–0.99
28	0.35	0.22–0.48	0.95	0.94–0.97	0.17	0.10–0.24	0.98	0.97–0.99
29	0.32	0.20–0.45	0.97	0.96–0.98	0.22	0.13–0.30	0.98	0.97–0.99
30	0.28	0.17–0.40	0.97	0.97–0.98	0.22	0.13–0.31	0.98	0.97–0.99
31	0.23	0.13–0.33	0.98	0.97–0.98	0.21	0.13–0.30	0.98	0.97–0.99
32	0.17	0.10–0.25	0.98	0.98–0.99	0.20	0.12–0.28	0.98	0.97–0.99
33	0.16	0.08–0.23	0.98	0.98–0.99	0.21	0.12–0.30	0.98	0.97–0.98
34	0.15	0.08–0.22	0.99	0.98–0.99	0.23	0.12–0.33	0.98	0.97–0.98
35	0.11	0.05–0.17	0.99	0.98–0.99	0.19	0.09–0.30	0.98	0.97–0.98
36	0.09	0.03–0.14	0.99	0.98–0.99	0.18	0.07–0.28	0.98	0.97–0.98
37	0.08	0.03–0.13	0.99	0.99–1.00	0.19	0.07–0.31	0.98	0.97–0.98
38	0.06	0.02–0.11	0.99	0.99–1.00	0.20	0.06–0.34	0.98	0.97–0.98
39	0.06	0.02–0.11	0.99	0.99–1.00	0.22	0.07–0.37	0.98	0.97–0.98
40	0.06	0.02–0.11	1.00	0.99–1.00	0.32	0.11–0.53	0.98	0.97–0.98
41	0.06	0.02–0.11	1.00	1.00–1.00	0.42	0.14–0.69	0.98	0.97–0.98
42	0.05	0.01–0.09	1.00	1.00–1.00	0.41	0.09–0.72	0.97	0.97–0.98
43	0.04	0.01–0.08	1.00	1.00–1.00	0.36	0.05–0.67	0.97	0.97–0.98
44	0.03	0.00–0.06	1.00	1.00–1.00	0.28	0.00–0.60	0.97	0.97–0.98

Cut-off	Sensitivity		Specificity		Positive predictive value		Negative predictive value	
	Value	95% CI	Value	95% CI	Value	95% CI	Value	95% CI
45	0.02	0.00–0.04	1.00	1.00–1.00	0.25	0.00–0.61	0.97	0.97–0.98
46	0.02	0.00–0.04	1.00	1.00–1.00	0.51	0.03–0.99	0.97	0.97–0.98
47	0.00	0.00–0.00	1.00	1.00–1.00	0.00	0.00–0.00	0.97	0.97–0.98
48	0.00	0.00–0.00	1.00	1.00–1.00	0.00	0.00–0.00	0.97	0.97–0.98
49	0.00	0.00–0.00	1.00	1.00–1.00	0.00	0.00–0.00	0.97	0.97–0.98
50	0.00	0.00–0.00	1.00	1.00–1.00	0.00	0.00–0.00	0.97	0.97–0.98

AUC	
Value	95% CI
0.81	0.70–0.91

Table B.28: Properties of the K10 for predicting 30-day ICD-10 anxiety or affective disorder

Cut-off	Sensitivity		Specificity		Positive predictive value		Negative predictive value	
	Value	95% CI	Value	95% CI	Value	95% CI	Value	95% CI
10	1.00	1.00–1.00	0.00	0.00–0.00	0.09	0.07–0.11	–	–
11	1.00	0.99–1.00	0.15	0.11–0.18	0.10	0.08–0.13	1.00	0.99–1.00
12	1.00	0.99–1.00	0.27	0.23–0.31	0.12	0.09–0.15	1.00	1.00–1.00
13	0.83	0.65–1.02	0.37	0.32–0.41	0.12	0.10–0.13	0.96	0.90–1.01
14	0.79	0.61–0.97	0.47	0.42–0.51	0.13	0.11–0.15	0.96	0.91–1.00
15	0.76	0.58–0.93	0.56	0.51–0.60	0.15	0.12–0.17	0.96	0.92–1.00
16	0.72	0.55–0.89	0.65	0.61–0.69	0.17	0.14–0.20	0.96	0.92–0.99
17	0.68	0.51–0.84	0.72	0.69–0.76	0.20	0.16–0.23	0.96	0.93–0.99
18	0.62	0.46–0.77	0.76	0.73–0.80	0.21	0.17–0.24	0.95	0.92–0.98
19	0.59	0.44–0.73	0.81	0.78–0.84	0.24	0.19–0.28	0.95	0.92–0.98
20	0.50	0.37–0.63	0.85	0.83–0.87	0.25	0.21–0.29	0.94	0.92–0.97
21	0.44	0.33–0.56	0.88	0.86–0.90	0.27	0.22–0.31	0.94	0.91–0.97
22	0.41	0.30–0.52	0.90	0.88–0.92	0.29	0.24–0.34	0.94	0.91–0.96
23	0.37	0.27–0.48	0.91	0.89–0.93	0.30	0.24–0.36	0.94	0.91–0.96
24	0.35	0.25–0.45	0.93	0.91–0.94	0.33	0.26–0.40	0.93	0.91–0.96
25	0.30	0.21–0.39	0.93	0.92–0.95	0.32	0.24–0.39	0.93	0.91–0.96
26	0.28	0.20–0.37	0.95	0.94–0.96	0.36	0.27–0.44	0.93	0.91–0.95
27	0.24	0.16–0.32	0.95	0.94–0.97	0.34	0.26–0.43	0.93	0.90–0.95
28	0.23	0.15–0.30	0.96	0.95–0.98	0.39	0.28–0.49	0.93	0.90–0.95
29	0.20	0.13–0.27	0.98	0.97–0.98	0.46	0.36–0.57	0.92	0.90–0.95
30	0.18	0.12–0.25	0.98	0.97–0.99	0.49	0.37–0.60	0.92	0.90–0.95
31	0.16	0.10–0.22	0.98	0.98–0.99	0.52	0.39–0.64	0.92	0.90–0.94
32	0.13	0.08–0.18	0.99	0.98–0.99	0.53	0.39–0.66	0.92	0.90–0.94
33	0.11	0.07–0.16	0.99	0.98–0.99	0.52	0.37–0.67	0.92	0.89–0.94
34	0.11	0.06–0.15	0.99	0.99–1.00	0.56	0.40–0.72	0.92	0.89–0.94
35	0.09	0.05–0.14	0.99	0.99–1.00	0.56	0.39–0.73	0.92	0.89–0.94
36	0.08	0.04–0.12	0.99	0.99–1.00	0.57	0.38–0.75	0.92	0.89–0.94
37	0.07	0.03–0.10	0.99	0.99–1.00	0.55	0.35–0.75	0.91	0.89–0.94
38	0.04	0.02–0.07	1.00	0.99–1.00	0.48	0.27–0.70	0.91	0.89–0.94
39	0.04	0.02–0.06	1.00	0.99–1.00	0.44	0.22–0.66	0.91	0.89–0.93
40	0.03	0.01–0.05	1.00	1.00–1.00	0.56	0.29–0.83	0.91	0.89–0.93
41	0.02	0.01–0.04	1.00	1.00–1.00	0.56	0.25–0.87	0.91	0.89–0.93
42	0.02	0.00–0.03	1.00	1.00–1.00	0.47	0.13–0.80	0.91	0.89–0.93
43	0.01	0.00–0.03	1.00	1.00–1.00	0.43	0.09–0.76	0.91	0.89–0.93
44	0.01	0.00–0.02	1.00	1.00–1.00	0.28	0.00–0.60	0.91	0.89–0.93

Cut-off	Sensitivity		Specificity		Positive predictive value		Negative predictive value	
	Value	95% CI	Value	95% CI	Value	95% CI	Value	95% CI
45	0.00	0.00–0.01	1.00	1.00–1.00	0.25	0.00–0.61	0.91	0.89–0.93
46	0.00	0.00–0.01	1.00	1.00–1.00	0.51	0.03–0.99	0.91	0.89–0.93
47	0.00	0.00–0.00	1.00	1.00–1.00	0.00	0.00–0.00	0.91	0.89–0.93
48	0.00	0.00–0.00	1.00	1.00–1.00	0.00	0.00–0.00	0.91	0.89–0.93
49	0.00	0.00–0.00	1.00	1.00–1.00	0.00	0.00–0.00	0.91	0.89–0.93
50	0.00	0.00–0.00	1.00	1.00–1.00	0.00	0.00–0.00	0.91	0.89–0.93

AUC	
Value	95% CI
0.75	0.63–0.86

Table B.29: PCL risk categories, by sex by Service, and by Service, rank and deployment status

	Total			Low (17-29)			Moderate (30-39)			High (40-49)			Very high (50-85)		
	N	%	% CI	N	%	% CI	N	%	% CI	N	%	% CI	N	%	% CI
Total	50,049	84.6	84.3-85.0	4,360	8.7	8.4-9.0	1,849	3.7	3.5-3.9	1,477	3.0	2.8-3.1	1,477	3.0	2.8-3.1
Males	43,241	84.7	84.3-85.2	3,749	8.7	8.3-9.0	1,578	3.7	3.4-3.9	1,269	2.9	2.7-3.1	1,269	2.9	2.7-3.1
Females	6,808	84.0	83.1-84.9	612	9.0	8.3-9.7	270	4.0	3.5-4.4	207	3.0	2.6-3.5	207	3.0	2.6-3.5
Navy	11,613	84.0	83.1-84.8	1,060	9.1	8.5-9.8	454	3.9	3.5-4.4	345	3.0	2.6-3.4	345	3.0	2.6-3.4
Males	9,508	84.4	83.4-85.3	846	8.9	8.2-9.6	351	3.7	3.2-4.2	289	3.0	2.6-3.5	289	3.0	2.6-3.5
Females	2,104	82.3	80.4-84.1	214	10.2	8.7-11.7	103	4.9	3.9-5.9	57	2.7	1.9-3.5	57	2.7	1.9-3.5
Army	25,356	83.5	82.9-84.1	2,327	9.2	8.7-9.6	1,006	4.0	3.7-4.3	850	3.4	3.1-3.6	850	3.4	3.1-3.6
Males	22,843	83.4	82.8-84.1	2,100	9.2	8.7-9.7	910	4.0	3.7-4.3	774	3.4	3.1-3.7	774	3.4	3.1-3.7
Females	2,513	84.1	82.7-85.5	227	9.0	7.9-10.2	96	3.8	3.1-4.6	75	3.0	2.3-3.7	75	3.0	2.3-3.7
Air Force	13,081	87.4	86.9-88.0	973	7.4	7.0-7.9	388	3.0	2.7-3.2	282	2.2	1.9-2.4	282	2.2	1.9-2.4
Males	10,890	87.8	87.2-88.4	803	7.4	6.9-7.8	317	2.9	2.6-3.2	206	1.9	1.7-2.1	206	1.9	1.7-2.1
Females	2,191	85.5	84.3-86.7	170	7.8	6.8-8.7	71	3.3	2.6-3.9	75	3.4	2.8-4.0	75	3.4	2.8-4.0
Officers	12,034	87.4	86.8-87.9	937	7.8	7.4-8.2	318	2.6	2.4-2.9	266	2.2	2.0-2.4	266	2.2	2.0-2.4
Non-commissioned officers	22,320	82.9	82.4-83.4	2,134	9.6	9.2-9.9	949	4.3	4.0-4.5	739	3.3	3.1-3.5	739	3.3	3.1-3.5
Other ranks	15,695	85.1	84.2-86.0	1,290	8.2	7.5-8.9	582	3.7	3.2-4.2	472	3.0	2.6-3.4	472	3.0	2.6-3.4
Ever deployed	30,850	84.1	83.7-84.6	2,815	9.1	8.8-9.5	1,140	3.7	3.5-3.9	941	3.0	2.8-3.3	941	3.0	2.8-3.3
Never deployed	19,199	85.5	84.8-86.1	1,545	8.0	7.5-8.6	708	3.7	3.3-4.0	536	2.8	2.5-3.1	536	2.8	2.5-3.1

Table B.30: Properties of the PCL for predicting 30-day ICD-10 post-traumatic stress disorder

Cut-off	Sensitivity		Specificity		Positive predictive value		Negative predictive value	
	Value	95% CI	Value	95% CI	Value	95% CI	Value	95% CI
17	1.00	1.00–1.00	0.00	0.00–0.00	0.03	0.02–0.04	–	–
18	1.00	0.99–1.00	0.33	0.29–0.37	0.05	0.04–0.06	1.00	1.00–1.00
19	0.96	0.89–1.00	0.40	0.36–0.44	0.05	0.04–0.07	1.00	0.99–1.00
20	0.96	0.89–1.00	0.48	0.44–0.52	0.06	0.04–0.07	1.00	0.99–1.00
21	0.93	0.85–1.00	0.55	0.51–0.59	0.07	0.05–0.08	1.00	0.99–1.00
22	0.91	0.83–1.00	0.59	0.55–0.63	0.07	0.05–0.09	1.00	0.99–1.00
23	0.86	0.74–0.98	0.61	0.58–0.65	0.07	0.05–0.09	0.99	0.98–1.00
24	0.80	0.67–0.93	0.65	0.61–0.68	0.07	0.05–0.09	0.99	0.98–1.00
25	0.80	0.67–0.93	0.67	0.64–0.71	0.08	0.06–0.10	0.99	0.98–1.00
26	0.80	0.67–0.93	0.72	0.70–0.75	0.09	0.07–0.11	0.99	0.98–1.00
27	0.80	0.67–0.93	0.75	0.72–0.77	0.10	0.07–0.12	0.99	0.98–1.00
28	0.79	0.65–0.92	0.78	0.75–0.80	0.11	0.08–0.13	0.99	0.98–1.00
29	0.79	0.65–0.92	0.80	0.77–0.82	0.12	0.09–0.15	0.99	0.98–1.00
30	0.74	0.60–0.87	0.82	0.80–0.84	0.12	0.09–0.15	0.99	0.98–1.00
31	0.74	0.60–0.87	0.83	0.82–0.85	0.13	0.10–0.16	0.99	0.98–1.00
32	0.69	0.56–0.83	0.85	0.83–0.87	0.14	0.10–0.17	0.99	0.98–0.99
33	0.66	0.52–0.79	0.86	0.85–0.88	0.14	0.11–0.18	0.99	0.98–0.99
34	0.64	0.51–0.78	0.88	0.86–0.89	0.15	0.11–0.19	0.99	0.98–0.99
35	0.63	0.50–0.76	0.89	0.87–0.90	0.16	0.12–0.20	0.99	0.98–0.99
36	0.63	0.49–0.76	0.89	0.88–0.91	0.17	0.13–0.21	0.99	0.98–0.99
37	0.61	0.48–0.74	0.91	0.89–0.92	0.18	0.14–0.23	0.99	0.98–0.99
38	0.53	0.41–0.66	0.91	0.90–0.93	0.18	0.13–0.22	0.98	0.98–0.99
39	0.50	0.37–0.62	0.92	0.91–0.93	0.18	0.13–0.23	0.98	0.97–0.99
40	0.47	0.35–0.60	0.93	0.92–0.94	0.18	0.13–0.23	0.98	0.97–0.99
41	0.46	0.34–0.58	0.93	0.92–0.94	0.19	0.13–0.24	0.98	0.97–0.99
42	0.43	0.31–0.55	0.94	0.93–0.95	0.19	0.14–0.25	0.98	0.97–0.99
43	0.42	0.30–0.54	0.94	0.93–0.95	0.20	0.14–0.25	0.98	0.97–0.99
44	0.40	0.29–0.52	0.94	0.93–0.95	0.20	0.14–0.26	0.98	0.97–0.99
45	0.38	0.27–0.50	0.95	0.94–0.96	0.21	0.14–0.27	0.98	0.97–0.99
46	0.36	0.25–0.47	0.95	0.94–0.96	0.21	0.14–0.27	0.98	0.97–0.99
47	0.33	0.23–0.44	0.96	0.95–0.96	0.21	0.14–0.28	0.98	0.97–0.98
48	0.32	0.21–0.43	0.96	0.95–0.97	0.22	0.15–0.29	0.98	0.97–0.98
49	0.30	0.19–0.40	0.96	0.95–0.97	0.21	0.14–0.29	0.98	0.97–0.98
50	0.30	0.19–0.40	0.97	0.96–0.97	0.23	0.15–0.31	0.98	0.97–0.98
51	0.29	0.19–0.40	0.97	0.96–0.98	0.24	0.16–0.33	0.98	0.97–0.98
52	0.28	0.18–0.39	0.97	0.96–0.98	0.26	0.17–0.35	0.98	0.97–0.98
53	0.25	0.15–0.35	0.97	0.97–0.98	0.26	0.16–0.36	0.97	0.97–0.98

Cut-off	Sensitivity		Specificity		Positive predictive value		Negative predictive value	
	Value	95% CI	Value	95% CI	Value	95% CI	Value	95% CI
54	0.21	0.12-0.30	0.98	0.97-0.98	0.24	0.14-0.34	0.97	0.96-0.98
55	0.20	0.12-0.29	0.98	0.97-0.98	0.24	0.14-0.34	0.97	0.96-0.98
56	0.20	0.12-0.29	0.98	0.97-0.99	0.26	0.15-0.37	0.97	0.96-0.98
57	0.17	0.10-0.25	0.98	0.98-0.99	0.25	0.14-0.35	0.97	0.96-0.98
58	0.16	0.09-0.24	0.98	0.98-0.99	0.25	0.14-0.37	0.97	0.96-0.98
59	0.16	0.09-0.24	0.98	0.98-0.99	0.26	0.14-0.38	0.97	0.96-0.98
60	0.16	0.09-0.24	0.99	0.98-0.99	0.28	0.15-0.40	0.97	0.96-0.98
61	0.15	0.08-0.23	0.99	0.98-0.99	0.29	0.15-0.43	0.97	0.96-0.98
62	0.13	0.05-0.20	0.99	0.98-0.99	0.27	0.13-0.42	0.97	0.96-0.98
63	0.13	0.05-0.20	0.99	0.99-1.00	0.33	0.16-0.51	0.97	0.96-0.98
64	0.11	0.04-0.18	0.99	0.99-1.00	0.33	0.15-0.51	0.97	0.96-0.98
65	0.11	0.04-0.18	0.99	0.99-1.00	0.36	0.17-0.56	0.97	0.96-0.98
66	0.10	0.03-0.17	0.99	0.99-1.00	0.39	0.18-0.60	0.97	0.96-0.98
67	0.08	0.02-0.14	1.00	0.99-1.00	0.44	0.20-0.68	0.97	0.96-0.98
68	0.08	0.02-0.14	1.00	1.00-1.00	0.49	0.24-0.74	0.97	0.96-0.98
69	0.04	0.01-0.08	1.00	1.00-1.00	0.37	0.14-0.60	0.97	0.96-0.98
70	0.04	0.01-0.08	1.00	1.00-1.00	0.39	0.15-0.63	0.97	0.96-0.98
71	0.03	0.00-0.05	1.00	1.00-1.00	0.35	0.07-0.63	0.97	0.96-0.98
72	0.01	0.00-0.02	1.00	1.00-1.00	0.21	0.05-0.48	0.97	0.96-0.98
73	0.01	0.00-0.02	1.00	1.00-1.00	0.26	0.06-0.58	0.97	0.96-0.98
74	0.01	0.00-0.02	1.00	1.00-1.00	0.39	0.06-0.83	0.97	0.96-0.98
75	0.01	0.00-0.02	1.00	1.00-1.00	0.39	0.06-0.83	0.97	0.96-0.98
76	0.01	0.00-0.02	1.00	1.00-1.00	0.39	0.06-0.83	0.97	0.96-0.98
77	0.01	0.00-0.02	1.00	1.00-1.00	0.39	0.06-0.83	0.97	0.96-0.98
78	0.01	0.00-0.01	1.00	1.00-1.00	0.25	0.19-0.68	0.97	0.96-0.98
79	0.01	0.00-0.01	1.00	1.00-1.00	0.25	0.19-0.68	0.97	0.96-0.98
80	0.01	0.00-0.01	1.00	1.00-1.00	0.25	0.19-0.68	0.97	0.96-0.98
81	0.01	0.00-0.01	1.00	1.00-1.00	0.25	0.19-0.68	0.97	0.96-0.98
82	0.00	0.00-0.00	1.00	1.00-1.00	0.00	0.00-0.00	0.97	0.96-0.98
83	0.00	0.00-0.00	1.00	1.00-1.00	0.00	0.00-0.00	0.97	0.96-0.98
84	0.00	0.00-0.00	1.00	1.00-1.00	0.00	0.00-0.00	0.97	0.96-0.98
85	0.00	0.00-0.00	1.00	1.00-1.00	0.00	0.00-0.00	0.97	0.96-0.98

AUC	
Value	95% CI
0.85	0.79-0.91

Table B.31: AUDIT risk categories, by sex by Service, and by Service, rank and deployment status

	N	Zone I (score 1-7)			Zone II (score 8-15)			Zone III (score 16-19)			Zone IV (score 20-40)		
		N	%	95% CI	N	%	95% CI	N	%	95% CI	N	%	95% CI
Total	50,049	36,848	73.6	73.1-74.2	11,345	22.7	22.2-23.2	1,172	2.3	2.1-2.5	684	1.4	1.2-1.5
Males	43,241	31,072	71.9	71.3-72.5	10,417	24.1	23.5-24.7	1,091	2.5	2.3-2.8	661	1.5	1.3-1.7
Navy	9,508	6,872	72.3	70.9-73.6	2,228	23.4	22.2-24.7	240	2.5	2.0-3.0	168	1.8	1.3-2.2
Army	22,843	15,428	67.5	66.6-68.5	6,251	27.4	26.5-28.3	720	3.2	2.8-3.5	444	1.9	1.7-2.2
Air Force	10,890	8,772	80.6	79.8-81.3	1,937	17.8	17.0-18.6	132	1.2	1.0-1.4	49	0.5	0.3-0.6
Females	6,808	5,776	84.8	83.9-85.8	928	13.6	12.7-14.6	81	1.2	0.9-1.5	23	0.3	0.2-0.5
Navy	2,104	1,721	81.8	79.7-83.9	346	16.4	14.4-18.5	23	1.1	0.5-1.7	14	0.6	0.2-1.1
Army	2,513	2,098	83.5	81.9-85.1	366	14.6	13.0-16.1	43	1.7	1.1-2.3	6	0.2	0.0-0.4
Air Force	2,191	1,956	89.3	88.1-90.5	217	9.9	8.7-11.0	15	0.7	0.4-1.0	3	0.1	0.0-0.4
Navy	11,612	8,594	74.0	72.9-75.2	2,574	22.2	21.1-23.3	263	2.3	1.8-2.7	182	1.6	1.2-1.9
Army	25,356	17,526	69.1	68.3-70.0	6,617	26.1	25.3-26.9	762	3.0	2.7-3.3	450	1.8	1.5-2.0
Air Force	13,081	10,728	82.0	81.3-82.7	2,154	16.5	15.8-17.1	147	1.1	0.9-1.3	52	0.4	0.3-0.5
Officers	12,034	9,667	80.3	79.7-81.0	2,103	17.5	16.8-18.1	204	1.7	1.5-1.9	59	0.5	0.4-0.6
Non-commissioned officers	22,319	16,878	75.6	75.0-76.3	4,774	21.4	20.8-22.0	408	1.8	1.6-2.0	260	1.2	1.0-1.3
Other ranks	15,696	10,303	65.6	64.3-67.0	4,468	28.5	27.2-29.8	561	3.6	3.0-4.1	365	2.3	1.9-2.8
Never deployed	19,010	13,877	73.0	72.0-74.0	4,394	23.1	22.2-24.0	459	2.4	2.0-2.8	280	1.5	1.2-1.8
Ever deployed	31,039	22,971	74.0	73.4-74.7	6,951	22.4	21.8-23.0	713	2.3	2.1-2.5	404	1.3	1.1-1.5

Table B.32: Frequency of alcohol consumption: How often do you have a drink containing alcohol?

	N	Never			Monthly or less			2 to 4 times a month			2 to 3 times a week			4 or more times a week		
		N	%	95% CI	N	%	95% CI	N	%	95% CI	N	%	95% CI	N	%	95% CI
Total	50,049	2,187	4.4	4.1-4.6	10,648	21.3	20.8-21.8	17,208	34.4	33.8-34.9	14,477	28.9	28.4-29.5	5,529	11.0	10.7-11.4
Males	43,241	1,774	4.1	3.8-4.4	8,565	19.8	19.3-20.3	14,951	34.6	33.9-35.2	12,952	30.0	29.4-30.5	4,999	11.6	11.2-11.9
Navy	9,508	328	3.4	2.9-4.0	1,947	20.5	19.3-21.7	3,335	35.1	33.7-36.5	2,920	30.7	29.4-32.0	978	10.3	9.5-11.1
Army	22,843	971	4.2	3.9-4.6	4,366	19.1	18.3-19.9	7,832	34.3	33.4-35.2	6,739	29.5	28.6-30.4	2,936	12.9	12.2-13.5
Air Force	10,890	476	4.4	4.0-4.8	2,251	20.7	19.9-21.5	3,785	34.8	33.8-35.7	3,293	30.2	29.4-31.1	1,085	10.0	9.4-10.5
Females	6,808	413	6.1	5.5-6.7	2,083	30.6	29.4-31.8	2,256	33.1	31.9-34.3	1,525	22.4	21.4-23.4	531	7.8	7.2-8.4
Navy	2,104	123	5.9	4.7-7.0	627	29.8	27.3-32.3	786	37.3	34.8-39.9	467	22.2	20.0-24.3	101	4.8	3.9-5.7
Army	2,513	152	6.0	5.1-7.0	777	30.9	29.0-32.9	784	31.2	29.3-33.1	540	21.5	19.8-23.2	260	10.4	9.2-11.5
Air Force	2,191	138	6.3	5.4-7.2	679	31.0	29.2-32.8	687	31.3	29.6-33.1	518	23.7	22.1-25.2	170	7.7	6.7-8.7
Navy	11,612	451	3.9	3.4-4.4	2,575	22.2	21.1-23.3	4,120	35.5	34.2-36.7	3,387	29.2	28.0-30.3	1,079	9.3	8.6-9.9
Army	25,356	1,123	4.4	4.1-4.8	5,143	20.3	19.6-21.0	8,616	34.0	33.1-34.8	7,279	28.7	27.9-29.5	3,196	12.6	12.0-13.2
Air Force	13,081	614	4.7	4.3-5.1	2,930	22.4	21.7-23.1	4,472	34.2	33.4-35.0	3,811	29.1	28.4-29.9	1,254	9.6	9.1-10.1
Officers	12,034	412	3.4	3.1-3.7	1,692	14.1	13.5-14.6	3,841	31.9	31.1-32.7	4,067	33.8	33.0-34.6	2,022	16.8	16.2-17.4
Non-commissioned officers	22,319	995	4.5	4.2-4.7	5,094	22.8	22.2-23.4	7,462	33.4	32.7-34.1	6,383	28.6	27.9-29.3	2,385	10.7	10.2-11.1
Other ranks	15,696	781	5.0	4.4-5.6	3,861	24.6	23.4-25.8	5,905	37.6	36.2-39.0	4,026	25.6	24.4-26.9	1,122	7.1	6.4-7.9
Never deployed	19,010	935	4.9	4.5-5.4	4,425	23.3	22.4-24.2	6,696	35.2	34.2-36.2	5,226	27.5	26.6-28.4	1,728	9.1	8.5-9.7
Ever deployed	31,040	1,253	4.0	3.8-4.3	6,223	20.0	19.5-20.6	10,511	33.9	33.2-34.5	9,251	29.8	29.2-30.5	3,802	12.2	11.8-12.7

Table B.33: Quantity of alcohol consumption: How many 'standard' drinks containing alcohol do you have on a typical day when you are drinking?

	N	N/A			1 or 2			3 or 4			5 or 6			7 or 9			10 or more		
		N	%	95% CI	N	%	95% CI	N	%	95% CI	N	%	95% CI	N	%	95% CI	N	%	95% CI
Total	50,049	2,877	5.7	5.5–6.0	15,504	31.0	30.5–31.5	17,127	34.2	33.7–34.8	8,684	17.4	16.9–17.8	3,439	6.9	6.5–7.2	2,419	4.8	4.5–5.1
Males	43,241	2,358	5.5	5.1–5.8	12,510	28.9	28.4–29.5	14,953	34.6	34.0–35.2	7,897	18.3	17.7–18.8	3,208	7.4	7.0–7.8	2,316	5.4	5.0–5.7
Navy	9,508	436	4.6	4.0–5.2	2,659	28.0	26.7–29.2	3,478	36.6	35.2–38.0	1,807	19.0	17.8–20.2	732	7.7	6.9–8.5	396	4.2	3.5–4.8
Army	22,843	1,312	5.7	5.3–6.2	6,086	26.6	25.9–27.4	7,494	32.8	31.9–33.7	4,391	19.2	18.4–20.0	1,916	8.4	7.8–9.0	1,643	7.2	6.6–7.8
Air Force	10,890	610	5.6	5.1–6.1	3,764	34.6	33.7–35.5	3,981	36.6	35.6–37.5	1,698	15.6	14.9–16.3	560	5.1	4.7–5.6	277	2.5	2.2–2.9
Females	6,808	519	7.6	7.0–8.3	2,995	44.0	42.8–45.2	2,174	31.9	30.7–33.1	786	11.5	10.7–12.4	231	3.4	2.9–3.9	103	1.5	1.2–1.9
Navy	2,104	144	6.9	5.6–8.1	809	38.4	36.0–40.9	711	33.8	31.3–36.4	323	15.3	13.4–17.3	89	4.3	3.1–5.4	27	1.3	0.7–1.9
Army	2,513	204	8.1	7.0–9.3	1,051	41.8	39.8–43.8	799	31.8	29.8–33.7	315	12.5	11.1–14.0	88	3.5	2.7–4.3	55	2.2	1.5–2.9
Air Force	2,191	170	7.8	6.7–8.8	1,134	51.8	49.9–53.7	664	30.3	28.6–32.1	148	6.7	5.8–7.7	54	2.5	1.8–3.1	21	0.9	0.5–1.4
Navy	11,612	580	5.0	4.4–5.5	3,468	29.9	28.8–31.0	4,190	36.1	34.9–37.3	2,130	18.3	17.3–19.4	822	7.1	6.4–7.8	423	3.6	3.1–4.2
Army	25,356	1,517	6.0	5.5–6.4	7,138	28.2	27.4–28.9	8,292	32.7	31.9–33.5	4,707	18.6	17.8–19.3	2,004	7.9	7.4–8.4	1,699	6.7	6.2–7.2
Air Force	13,081	780	6.0	5.5–6.4	4,899	37.5	36.6–38.3	4,645	35.5	34.7–36.3	1,846	14.1	13.5–14.7	614	4.7	4.3–5.1	297	2.3	2.0–2.6
Officers	12,034	511	4.2	3.9–4.6	5,253	43.6	42.8–44.5	4,258	35.4	34.6–36.2	1,386	11.5	11.0–12.1	464	3.9	3.5–4.2	163	1.4	1.1–1.6
Non-commissioned officers	22,319	1,318	5.9	5.6–6.2	7,032	31.5	30.8–32.2	7,917	35.5	34.8–36.2	3,827	17.1	16.6–17.7	1,382	6.2	5.8–6.6	842	3.8	3.5–4.1
Other ranks	15,696	1,047	6.7	6.0–7.4	3,219	20.5	19.4–21.6	4,952	31.6	30.2–32.9	3,470	22.1	20.9–23.3	1,594	10.2	9.3–11.0	1,414	9.0	8.2–9.9
Never deployed	19,007	1,189	6.3	5.8–6.8	5,546	29.2	28.3–30.0	6,271	33.0	32.0–33.9	3,517	18.5	17.7–19.3	1,473	7.7	7.1–8.4	1,014	5.3	4.8–5.9
Ever deployed	31,043	1,688	5.4	5.1–5.8	9,958	32.1	31.5–32.7	10,856	35.0	34.3–35.7	5,166	16.6	16.1–17.2	1,967	6.3	6.0–6.7	1,405	4.5	4.2–4.9

Table B.34: Percentage of ADF reporting a problem with drinking

	N	No			Probably not			Unsure			Possibly			Definitely		
		N	%	95% CI	N	%	95% CI	N	%	95% CI	N	%	95% CI	N	%	95% CI
Total	50,049	45,776	91.5	91.1–91.8	2,547	5.1	4.8–5.4	713	1.4	1.3–1.6	827	1.7	1.5–1.8	186	0.4	0.3–0.4
Males	43,241	39,262	90.8	90.4–91.2	2,369	5.5	5.2–5.8	657	1.5	1.3–1.7	768	1.8	1.6–1.9	184	0.4	0.3–0.5
Navy	9,508	8,531	89.7	88.8–90.6	558	5.9	5.2–6.5	187	2.0	1.5–2.4	186	2.0	1.6–2.4	46	0.5	0.3–0.7
Army	22,843	20,508	89.8	89.2–90.4	1,365	6.0	5.5–6.4	365	1.6	1.3–1.9	495	2.2	1.9–2.4	111	0.5	0.4–0.6
Air Force	10,890	10,224	93.9	93.4–94.3	447	4.1	3.7–4.5	106	1.0	0.8–1.2	87	0.8	0.6–1.0	27	0.3	0.1–0.4
Females	6,808	6,513	95.7	95.2–96.2	178	2.6	2.2–3.0	55	0.8	0.6–1.1	60	0.9	0.7–1.1	2	0.0	0.0–0.1
Navy	2,104	2,023	96.2	95.3–97.1	49	2.3	1.7–2.9	21	1.0	0.4–1.6	10	0.5	0.2–0.8	0	0.0	0.0–0.0
Army	2,513	2,380	94.7	93.8–95.6	71	2.8	2.2–3.5	29	1.2	0.7–1.6	31	1.2	0.8–1.6	2	0.1	0.0–0.2
Air Force	2,191	2,110	96.3	95.6–97.0	58	2.6	2.0–3.3	5	0.2	0.1–0.4	18	0.8	0.5–1.2	0	0.0	0.0–0.0
Navy	11,612	10,554	90.9	90.1–91.6	607	5.2	4.7–5.8	208	1.8	1.4–2.2	196	1.7	1.4–2.0	46	0.4	0.3–0.5
Army	25,356	22,888	90.3	89.7–90.8	1,436	5.7	5.2–6.1	394	1.6	1.3–1.8	526	2.1	1.8–2.3	113	0.4	0.3–0.6
Air Force	13,081	12,334	94.3	93.9–94.7	504	3.9	3.5–4.2	111	0.8	0.7–1.0	105	0.8	0.7–0.9	27	0.2	0.1–0.3
Officers	12,034	11,073	92.0	91.6–92.5	614	5.1	4.7–5.5	124	1.0	0.9–1.2	196	1.6	1.4–1.8	28	0.2	0.1–0.3
Non-commissioned officers	22,319	20,401	91.4	91.0–91.8	1,134	5.1	4.8–5.4	274	1.2	1.1–1.4	414	1.9	1.7–2.1	97	0.4	0.3–0.5
Other ranks	15,696	14,302	91.1	90.3–91.9	799	5.1	4.5–5.7	315	2.0	1.6–2.4	218	1.4	1.0–1.7	62	0.4	0.2–0.6
Never deployed	19,010	17,516	92.1	91.6–92.7	942	5.0	4.5–5.4	251	1.3	1.0–1.6	257	1.4	1.1–1.6	44	0.2	0.1–0.3
Ever deployed	31,040	28,260	91.0	90.6–91.5	1,605	5.2	4.9–5.5	462	1.5	1.3–1.7	571	1.8	1.7–2.0	142	0.5	0.4–0.6

Table B.35: Percentage of ADF who reported difficulties reducing their alcohol intake

	N	Very easy			Fairly easy			Neither difficult nor easy			Fairly difficult			Very difficult			N/A		
		N	%	95% CI	N	%	95% CI	N	%	95% CI	N	%	95% CI	N	%	95% CI	N	%	
Total	50,049	32,543	65.0	64.5–65.6	8,296	16.6	16.1–17.0	3,228	6.4	6.2–6.8	1,057	2.1	1.9–2.3	213	0.4	0.3–0.5	4,711	9.4	9.1–9.7
Males	43,241	27,821	64.3	63.7–65.0	7,505	17.4	16.9–17.9	2,982	6.9	6.6–7.2	961	2.2	2.0–2.4	209	0.5	0.4–0.6	3,762	8.7	8.3–9.1
Navy	9,508	6,145	64.6	63.2–66.0	1,718	18.1	16.9–19.2	667	7.0	6.2–7.8	261	2.7	2.3–3.2	45	0.5	0.3–0.7	672	7.1	6.3–7.8
Army	22,843	14,571	63.8	62.9–64.7	3,937	17.2	16.5–18.0	1,685	7.4	6.9–7.9	519	2.3	2.0–2.6	132	0.6	0.4–0.7	1,999	8.8	8.2–9.3
Air Force	10,890	7,105	65.2	64.3–66.2	1,850	17.0	16.3–17.7	631	5.8	5.3–6.3	181	1.7	1.4–1.9	32	0.3	0.2–0.4	1,090	10.0	9.4–10.6
Females	6,808	4,722	69.4	68.2–70.5	791	11.6	10.8–12.4	246	3.6	3.1–4.1	95	1.4	1.1–1.7	4	0.1	0.0–0.1	950	13.9	13.1–14.8
Navy	2,104	1,542	73.3	71.0–75.6	219	10.4	8.8–12.0	72	3.4	2.5–4.3	26	1.2	0.7–1.8	4	0.2	0.0–0.4	242	11.5	9.8–13.2
Army	2,513	1,666	66.3	64.3–68.3	327	13.0	11.6–14.4	103	4.1	3.3–4.9	29	1.1	0.7–1.5	0	0.0	0.0–0.0	388	15.5	14.0–16.9
Air Force	2,191	1,514	69.1	67.3–70.9	245	11.2	10.0–12.4	71	3.3	2.6–3.9	41	1.9	1.4–2.4	0	0.0	0.0–0.0	319	14.6	13.2–15.9
Navy	11,612	7,687	66.2	65.0–67.4	1,937	16.7	15.7–17.6	738	6.4	5.7–7.0	287	2.5	2.1–2.9	49	0.4	0.2–0.6	914	7.9	7.2–8.6
Army	25,356	16,237	64.0	63.2–64.9	4,264	16.8	16.1–17.5	1,788	7.1	6.6–7.5	547	2.2	1.9–2.4	132	0.5	0.4–0.7	2,388	9.4	8.9–9.9
Air Force	13,081	8,619	65.9	65.1–66.7	2,096	16.0	15.4–16.7	703	5.4	5.0–5.8	222	1.7	1.5–1.9	32	0.2	0.2–0.3	1,409	10.8	10.2–11.3
Officers	12,034	7,984	66.3	65.6–67.1	2,174	18.1	17.4–18.7	714	5.9	5.5–6.3	266	2.2	2.0–2.5	40	0.3	0.2–0.4	855	7.1	6.7–7.5
Non-commissioned officers	22,319	14,500	65.0	64.3–65.7	3,484	15.6	15.1–16.1	1,431	6.4	6.0–6.8	458	2.1	1.8–2.3	76	0.3	0.3–0.4	2,370	10.6	10.2–11.1
Other ranks	15,696	10,058	64.1	62.7–65.5	2,638	16.8	15.7–17.9	1,084	6.9	6.2–7.6	333	2.1	1.7–2.6	97	0.6	0.4–0.9	1,486	9.5	8.7–10.3
Never deployed	19,009	11,920	62.7	61.7–63.7	3,169	16.7	15.9–17.4	1,259	6.6	6.1–7.2	445	2.3	2.0–2.7	78	0.4	0.3–0.6	2,138	11.2	10.6–11.9
Ever deployed	31,039	20,622	66.4	65.8–67.1	5,127	16.5	16.0–17.1	1,970	6.3	6.0–6.7	612	2.0	1.8–2.2	135	0.4	0.3–0.5	2,573	8.3	7.9–8.7

Table B.36: Properties of AUDIT for predicting 30-day ICD-10 any alcohol disorder

Cut-off	Sensitivity		Specificity		Positive predictive value		Negative predictive value	
	Value	95% CI	Value	95% CI	Value	95% CI	Value	95% CI
0	1.00	1.00–1.00	0.00	0.00–0.00	0.01	0.01–0.01	–	–
1	1.00	1.00–1.00	0.05	0.02–0.07	0.01	0.01–0.02	1.00	1.00–1.00
2	1.00	1.00–1.00	0.13	0.10–0.16	0.01	0.01–0.02	1.00	1.00–1.00
3	1.00	1.00–1.00	0.21	0.18–0.25	0.01	0.01–0.02	1.00	1.00–1.00
4	1.00	1.00–1.00	0.32	0.28–0.36	0.01	0.01–0.02	1.00	1.00–1.00
5	1.00	1.00–1.00	0.49	0.45–0.53	0.02	0.01–0.03	1.00	1.00–1.00
6	1.00	1.00–1.00	0.60	0.56–0.63	0.02	0.01–0.04	1.00	1.00–1.00
7	1.00	1.00–1.00	0.70	0.66–0.73	0.03	0.02–0.05	1.00	1.00–1.00
8	0.95	0.89–1.00	0.76	0.73–0.78	0.04	0.02–0.06	1.00	1.00–1.00
9	0.83	0.64–1.00	0.83	0.81–0.85	0.05	0.02–0.07	1.00	1.00–1.00
10	0.72	0.50–0.94	0.86	0.84–0.88	0.05	0.02–0.07	1.00	0.99–1.00
11	0.72	0.50–0.94	0.88	0.87–0.90	0.06	0.03–0.09	1.00	0.99–1.00
12	0.63	0.38–0.87	0.91	0.90–0.92	0.07	0.03–0.10	1.00	0.99–1.00
13	0.49	0.25–0.72	0.94	0.93–0.95	0.07	0.03–0.11	0.99	0.99–1.00
14	0.43	0.20–0.65	0.95	0.94–0.96	0.08	0.03–0.13	0.99	0.99–1.00
15	0.41	0.18–0.64	0.96	0.95–0.97	0.10	0.03–0.16	0.99	0.99–1.00
16	0.38	0.16–0.61	0.97	0.96–0.98	0.12	0.04–0.20	0.99	0.99–1.00
17	0.28	0.09–0.47	0.98	0.97–0.99	0.13	0.04–0.22	0.99	0.99–1.00
18	0.24	0.06–0.43	0.99	0.98–0.99	0.16	0.04–0.28	0.99	0.99–1.00
19	0.21	0.03–0.39	0.99	0.99–0.99	0.17	0.03–0.32	0.99	0.99–1.00
20	0.19	0.02–0.37	0.99	0.99–1.00	0.22	0.03–0.41	0.99	0.99–1.00
21	0.08	0.00–0.16	0.99	0.99–1.00	0.12	0.00–0.23	0.99	0.99–1.00
22	0.08	0.00–0.16	1.00	0.99–1.00	0.14	0.00–0.29	0.99	0.99–1.00
23	0.06	0.00–0.13	1.00	0.99–1.00	0.14	0.00–0.32	0.99	0.99–1.00
24	0.05	0.00–0.11	1.00	0.99–1.00	0.14	0.00–0.35	0.99	0.99–1.00
25	0.05	0.00–0.11	1.00	0.99–1.00	0.14	0.00–0.35	0.99	0.99–1.00
26	0.03	0.00–0.08	1.00	1.00–1.00	0.13	0.00–0.38	0.99	0.99–1.00
27	0.03	0.00–0.08	1.00	1.00–1.00	0.29	0.00–0.75	0.99	0.99–1.00
28	0.00	0.00–0.00	1.00	1.00–1.00	0.00	0.00–0.00	0.99	0.99–0.99
29	0.00	0.00–0.00	1.00	1.00–1.00	0.00	0.00–0.00	0.99	0.99–0.99
30	0.00	0.00–0.00	1.00	1.00–1.00	–	–	0.99	0.99–0.99
31	0.00	0.00–0.00	1.00	1.00–1.00	–	–	0.99	0.99–0.99
32	0.00	0.00–0.00	1.00	1.00–1.00	–	–	0.99	0.99–0.99
33	0.00	0.00–0.00	1.00	1.00–1.00	–	–	0.99	0.99–0.99
34	0.00	0.00–0.00	1.00	1.00–1.00	–	–	0.99	0.99–0.99

Cut-off	Sensitivity		Specificity		Positive predictive value		Negative predictive value	
	Value	95% CI	Value	95% CI	Value	95% CI	Value	95% CI
35	0.00	0.00–0.00	1.00	1.00–1.00	–	–	0.99	0.99–0.99
36	0.00	0.00–0.00	1.00	1.00–1.00	–	–	0.99	0.99–0.99
37	0.00	0.00–0.00	1.00	1.00–1.00	–	–	0.99	0.99–0.99
38	0.00	0.00–0.00	1.00	1.00–1.00	–	–	0.99	0.99–0.99
39	0.00	0.00–0.00	1.00	1.00–1.00	–	–	0.99	0.99–0.99
40	0.00	0.00–0.00	1.00	1.00–1.00	–	–	0.99	0.99–0.99

AUC	
Value	95% CI
0.91	0.87–0.96

Table B.37: Properties of AUDIT for predicting 30-day ICD-10 alcohol harmful use

Cut-off	Sensitivity		Specificity		Positive predictive value		Negative predictive value	
	Value	95% CI	Value	95% CI	Value	95% CI	Value	95% CI
0	1.00	1.00–1.00	0.00	0.00–0.00	0.00	0.00–0.00	–	–
1	1.00	1.00–1.00	0.05	0.02–0.07	0.00	0.00–0.00	1.00	1.00–1.00
2	1.00	1.00–1.00	0.13	0.09–0.16	0.00	0.00–0.01	1.00	1.00–1.00
3	1.00	1.00–1.00	0.21	0.17–0.25	0.00	0.00–0.01	1.00	1.00–1.00
4	1.00	1.00–1.00	0.32	0.28–0.36	0.00	0.00–0.01	1.00	1.00–1.00
5	1.00	1.00–1.00	0.49	0.45–0.53	0.00	0.00–0.01	1.00	1.00–1.00
6	1.00	1.00–1.00	0.59	0.55–0.63	0.01	0.00–0.01	1.00	1.00–1.00
7	1.00	1.00–1.00	0.69	0.66–0.72	0.01	0.00–0.01	1.00	1.00–1.00
8	1.00	1.00–1.00	0.75	0.73–0.78	0.01	0.00–0.02	1.00	1.00–1.00
9	0.57	0.07–1.00	0.82	0.80–0.84	0.01	0.00–0.01	1.00	1.00–1.00
10	0.47	0.03–0.92	0.85	0.84–0.87	0.01	0.00–0.01	1.00	1.00–1.00
11	0.47	0.03–0.92	0.88	0.86–0.89	0.01	0.00–0.02	1.00	1.00–1.00
12	0.47	0.03–0.92	0.91	0.89–0.92	0.01	0.00–0.02	1.00	1.00–1.00
13	0.40	0.00–0.80	0.93	0.92–0.94	0.01	0.00–0.03	1.00	1.00–1.00
14	0.15	0.07–0.36	0.95	0.94–0.96	0.01	0.00–0.02	1.00	1.00–1.00
15	0.15	0.07–0.36	0.96	0.95–0.97	0.01	0.00–0.02	1.00	1.00–1.00
16	0.15	0.07–0.36	0.97	0.96–0.98	0.01	0.00–0.03	1.00	1.00–1.00
17	0.15	0.07–0.36	0.98	0.97–0.98	0.02	0.01–0.04	1.00	1.00–1.00
18	0.15	0.07–0.36	0.99	0.98–0.99	0.02	0.01–0.05	1.00	1.00–1.00
19	0.15	0.07–0.36	0.99	0.98–0.99	0.03	0.01–0.07	1.00	1.00–1.00
20	0.08	0.08–0.24	0.99	0.99–1.00	0.02	0.02–0.06	1.00	1.00–1.00
21	0.08	0.08–0.24	0.99	0.99–1.00	0.03	0.03–0.08	1.00	1.00–1.00
22	0.08	0.08–0.24	0.99	0.99–1.00	0.03	0.03–0.10	1.00	1.00–1.00
23	0.00	0.00–0.00	1.00	0.99–1.00	0.00	0.00–0.00	1.00	1.00–1.00
24	0.00	0.00–0.00	1.00	0.99–1.00	0.00	0.00–0.00	1.00	1.00–1.00
25	0.00	0.00–0.00	1.00	0.99–1.00	0.00	0.00–0.00	1.00	1.00–1.00
26	0.00	0.00–0.00	1.00	1.00–1.00	0.00	0.00–0.00	1.00	1.00–1.00
27	0.00	0.00–0.00	1.00	1.00–1.00	0.00	0.00–0.00	1.00	1.00–1.00
28	0.00	0.00–0.00	1.00	1.00–1.00	0.00	0.00–0.00	1.00	1.00–1.00
29	0.00	0.00–0.00	1.00	1.00–1.00	0.00	0.00–0.00	1.00	1.00–1.00
30	0.00	0.00–0.00	1.00	1.00–1.00	–	–	1.00	1.00–1.00
31	0.00	0.00–0.00	1.00	1.00–1.00	–	–	1.00	1.00–1.00
32	0.00	0.00–0.00	1.00	1.00–1.00	–	–	1.00	1.00–1.00
33	0.00	0.00–0.00	1.00	1.00–1.00	–	–	1.00	1.00–1.00
34	0.00	0.00–0.00	1.00	1.00–1.00	–	–	1.00	1.00–1.00

Cut-off	Sensitivity		Specificity		Positive predictive value		Negative predictive value	
	Value	95% CI	Value	95% CI	Value	95% CI	Value	95% CI
35	0.00	0.00–0.00	1.00	1.00–1.00	–	–	1.00	1.00–1.00
36	0.00	0.00–0.00	1.00	1.00–1.00	–	–	1.00	1.00–1.00
37	0.00	0.00–0.00	1.00	1.00–1.00	–	–	1.00	1.00–1.00
38	0.00	0.00–0.00	1.00	1.00–1.00	–	–	1.00	1.00–1.00
39	0.00	0.00–0.00	1.00	1.00–1.00	–	–	1.00	1.00–1.00
40	0.00	0.00–0.00	1.00	1.00–1.00	–	–	1.00	1.00–1.00

AUC	
Value	95% CI
0.87	0.72–0.98

Table B.38: Properties of AUDIT for predicting 30-day ICD-10 alcohol dependence

Cut-off	Sensitivity		Specificity		Positive predictive value		Negative predictive value	
	Value	95% CI	Value	95% CI	Value	95% CI	Value	95% CI
0	1.00	1.00–1.00	0.00	0.00–0.00	0.01	0.00–0.01	–	–
1	1.00	1.00–1.00	0.05	0.02–0.07	0.01	0.00–0.01	1.00	1.00–1.00
2	1.00	1.00–1.00	0.13	0.10–0.16	0.01	0.00–0.01	1.00	1.00–1.00
3	1.00	1.00–1.00	0.21	0.18–0.25	0.01	0.00–0.01	1.00	1.00–1.00
4	1.00	1.00–1.00	0.32	0.28–0.36	0.01	0.00–0.02	1.00	1.00–1.00
5	1.00	1.00–1.00	0.49	0.45–0.53	0.01	0.01–0.02	1.00	1.00–1.00
6	1.00	1.00–1.00	0.59	0.56–0.63	0.02	0.01–0.03	1.00	1.00–1.00
7	1.00	1.00–1.00	0.69	0.66–0.73	0.02	0.01–0.04	1.00	1.00–1.00
8	0.94	0.85–1.00	0.76	0.73–0.78	0.03	0.01–0.04	1.00	1.00–1.00
9	0.91	0.81–1.00	0.83	0.81–0.85	0.04	0.02–0.06	1.00	1.00–1.00
10	0.80	0.59–1.00	0.86	0.84–0.88	0.04	0.02–0.07	1.00	1.00–1.00
11	0.80	0.59–1.00	0.88	0.87–0.90	0.05	0.02–0.08	1.00	1.00–1.00
12	0.67	0.41–0.94	0.91	0.90–0.92	0.05	0.02–0.09	1.00	0.99–1.00
13	0.51	0.24–0.79	0.94	0.92–0.95	0.06	0.02–0.10	1.00	0.99–1.00
14	0.51	0.24–0.79	0.95	0.94–0.96	0.07	0.02–0.12	1.00	0.99–1.00
15	0.49	0.22–0.76	0.96	0.95–0.97	0.09	0.03–0.15	1.00	0.99–1.00
16	0.45	0.18–0.72	0.97	0.96–0.98	0.11	0.03–0.19	1.00	0.99–1.00
17	0.33	0.09–0.56	0.98	0.97–0.99	0.11	0.03–0.20	0.99	0.99–1.00
18	0.27	0.05–0.50	0.99	0.98–0.99	0.14	0.02–0.26	0.99	0.99–1.00
19	0.23	0.01–0.45	0.99	0.99–0.99	0.14	0.00–0.29	0.99	0.99–1.00
20	0.23	0.01–0.45	0.99	0.99–1.00	0.20	0.01–0.39	0.99	0.99–1.00
21	0.08	0.01–0.18	0.99	0.99–1.00	0.09	0.02–0.19	0.99	0.99–1.00
22	0.08	0.01–0.18	0.99	0.99–1.00	0.11	0.02–0.23	0.99	0.99–1.00
23	0.08	0.01–0.18	1.00	0.99–1.00	0.14	0.03–0.32	0.99	0.99–1.00
24	0.06	0.03–0.15	1.00	0.99–1.00	0.14	0.06–0.35	0.99	0.99–1.00
25	0.06	0.03–0.15	1.00	0.99–1.00	0.14	0.06–0.35	0.99	0.99–1.00
26	0.04	0.03–0.11	1.00	1.00–1.00	0.13	0.12–0.38	0.99	0.99–1.00
27	0.04	0.03–0.11	1.00	1.00–1.00	0.29	0.17–0.75	0.99	0.99–1.00
28	0.00	0.00–0.00	1.00	1.00–1.00	0.00	0.00–0.00	0.99	0.99–1.00
29	0.00	0.00–0.00	1.00	1.00–1.00	0.00	0.00–0.00	0.99	0.99–1.00
30	0.00	0.00–0.00	1.00	1.00–1.00	–	–	0.99	0.99–1.00
31	0.00	0.00–0.00	1.00	1.00–1.00	–	–	0.99	0.99–1.00
32	0.00	0.00–0.00	1.00	1.00–1.00	–	–	0.99	0.99–1.00
33	0.00	0.00–0.00	1.00	1.00–1.00	–	–	0.99	0.99–1.00
34	0.00	0.00–0.00	1.00	1.00–1.00	–	–	0.99	0.99–1.00

Cut-off	Sensitivity		Specificity		Positive predictive value		Negative predictive value	
	Value	95% CI	Value	95% CI	Value	95% CI	Value	95% CI
35	0.00	0.00–0.00	1.00	1.00–1.00	–	–	0.99	0.99–1.00
36	0.00	0.00–0.00	1.00	1.00–1.00	–	–	0.99	0.99–1.00
37	0.00	0.00–0.00	1.00	1.00–1.00	–	–	0.99	0.99–1.00
38	0.00	0.00–0.00	1.00	1.00–1.00	–	–	0.99	0.99–1.00
39	0.00	0.00–0.00	1.00	1.00–1.00	–	–	0.99	0.99–1.00
40	0.00	0.00–0.00	1.00	1.00–1.00	–	–	0.99	0.99–1.00

AUC	
Value	95% CI
0.93	0.89–0.97

Section 3 data tables

Table B.39: Estimated prevalence of reported stigma and barriers to care in the non-MEAO subpopulation, by type and sex

	Males			Females			Persons		
	N=26,169			N=4,679			N=30,848		
	N	%	95% CI	N	%	95% CI	N	%	95% CI
Q5.1 Proportion that sought help	4,190	16.01	(15.4, 16.7)	1,332	28.5	(27.0, 29.9)	5,522	17.9	(17.3, 18.5)
Q5.2 Not knowing where to get help									
Strongly disagree	11,928	45.6	(44.7, 46.5)	2,450	52.4	(50.7, 54.0)	14,378	46.6	(45.8, 47.4)
Disagree	10,413	39.8	(38.9, 40.7)	1,711	36.6	(35.0, 38.1)	12,124	39.3	(38.5, 40.1)
Uncertain	2,140	8.2	(7.6, 8.7)	263	5.6	(4.8, 6.4)	2,403	7.8	(7.3, 8.3)
Agree	1,379	5.3	(4.8, 5.7)	213	4.6	(3.8, 5.3)	1,592	5.2	(4.8, 5.5)
Strongly agree	310	1.2	(1.0, 1.4)	41	0.9	(0.6, 1.2)	351	1.1	(1.0, 1.3)
Q5.3 Difficulty getting time off work									
Strongly disagree	9,641	36.8	(36.0, 37.7)	1,756	37.5	(36.0, 39.1)	11,396	36.9	(36.2, 37.7)
Disagree	9,365	35.8	(34.9, 36.7)	1,596	34.1	(32.6, 35.7)	10,960	35.5	(34.8, 36.3)
Uncertain	3,311	12.7	(12.0, 13.3)	651	13.9	(12.8, 15.1)	3,962	12.8	(12.3, 13.4)
Agree	2,773	10.6	(10.0, 11.2)	523	11.2	(10.1, 12.2)	3,296	10.7	(10.2, 11.2)
Strongly agree	1,080	4.1	(3.7, 4.5)	153	3.3	(2.7, 3.9)	1,233	4.0	(3.6, 4.3)
Q5.4 Harm my career or career prospects									
Strongly disagree	6,477	24.7	(24.0, 25.5)	1,203	25.7	(24.3, 27.1)	7,680	24.9	(24.2, 25.6)
Disagree	6,847	26.2	(25.4, 27.0)	1,244	26.6	(25.2, 28.0)	8,091	26.2	(25.5, 26.9)
Uncertain	5,813	22.2	(21.4, 23.0)	958	20.5	(19.1, 21.8)	6,771	22.0	(21.3, 22.6)
Agree	4,739	18.1	(17.4, 18.8)	848	18.1	(16.8, 19.4)	5,587	18.1	(17.5, 18.7)
Strongly agree	2,293	8.8	(8.2, 9.3)	426	9.1	(8.1, 10.1)	2,719	8.8	(8.3, 9.3)

	Males			Females			Persons		
	N=26,169			N=4,679			N=30,848		
	N	%	95% CI	N	%	95% CI	N	%	95% CI
Q5.5 People would treat me differently									
Strongly disagree	5,900	22.5	(21.8,23.3)	1,119	23.9	(22.5,25.3)	7,019	22.8	(22.1,23.4)
Disagree	6,859	26.2	(25.4,27.0)	1,260	26.9	(25.5,28.4)	8,118	26.3	(25.6,27.0)
Uncertain	6,197	23.7	(22.9,24.5)	1,001	21.4	(20.0,22.8)	7,198	23.3	(22.6,24.0)
Agree	5,450	20.8	(20.1,21.6)	998	21.3	(20.0,22.7)	6,447	20.9	(20.2,21.6)
Strongly agree	1,764	6.7	(6.3,7.2)	302	6.4	(5.6,7.3)	2,065	6.7	(6.3,7.1)
Q5.6 Seen as weak									
Strongly disagree	6,408	24.5	(23.7,25.3)	1,217	26.0	(24.6,27.4)	7,625	24.7	(24.0,25.4)
Disagree	7,913	30.2	(29.4,31.1)	1,393	29.8	(28.3,31.3)	9,306	30.2	(29.4,30.9)
Uncertain	5,255	20.1	(19.3,20.8)	871	18.6	(17.3,19.9)	6,126	19.9	(19.2,20.5)
Agree	4,717	18.0	(17.3,18.7)	872	18.6	(17.4,19.9)	5,589	18.1	(17.5,18.7)
Strongly agree	1,877	7.2	(6.7,7.7)	326	7.0	(6.1,7.8)	2,202	7.1	(6.7,7.6)
Q5.7 Stop me from being deployed									
Strongly disagree	4,889	18.7	(18.0,19.4)	1,015	21.7	(20.3,23.0)	5,904	19.1	(18.5,19.8)
Disagree	4,518	17.3	(16.6,18.0)	811	17.3	(16.1,18.6)	5,329	17.3	(16.7,17.9)
Uncertain	7,071	27.0	(26.2,27.8)	1,169	25.0	(23.6,26.4)	8,240	26.7	(26.0,27.4)
Agree	5,974	22.8	(22.1,23.6)	1,052	22.5	(21.1,23.8)	7,025	22.8	(22.1,23.5)
Strongly agree	3,718	14.2	(13.6,14.9)	633	13.5	(12.4,14.7)	4,350	14.1	(13.5,14.7)

Table B.40: Prevalence of reported stigma and barriers to care in the ADF, by type and rank

	Officers			Non-commissioned officers			Other ranks		
	N=7,017			N=11,188			N=12,643		
	N	%	95% CI	N	%	95% CI	N	%	95% CI
Q5.1 Proportion that sought help	1,126	16	(15.2,16.9)	2,042	18.3	(17.4, 19.1)	2354	18.6	(17.5, 19.8)
Q5.2 Not knowing where to get help									
Strongly disagree	3,687	52.5	(51.3, 53.8)	5,139	45.9	(44.8,47.0)	5,552	43.9	(42.3, 45.5)
Disagree	2,729	38.9	(37.7, 40.1)	4,639	41.7	(40.4, 42.5)	4,756	37.6	(36.0, 39.2)
Uncertain	306	4.4	(3.9, 4.9)	733	6.6	(6.0, 7.1)	1,364	10.8	(9.8, 11.8)
Agree	233	3.3	(2.9, 3.8)	553	4.9	(4.5, 5.4)	806	6.4	(5.6, 7.1)
Strongly agree	62	0.9	(0.7, 1.1)	124	1.1	(0.9, 1.3)	165	1.3	(0.9, 1.7)
Q5.3 Difficulty getting time off work									
Strongly disagree	3,110	44.3	(43.1,45.5)	4,365	39.0	(38.0, 40.1)	3,921	31	(29.5, 32.5)
Disagree	2,501	35.6	(34.5, 36.8)	4,285	38.3	(37.2, 39.4)	4,174	33	(31.5, 34.5)
Uncertain	561	8.0	(7.3, 8.7)	1,246	11.1	(10.4, 11.8)	2,155	17	(15.8, 18.3)
Agree	677	9.6	(8.9, 10.4)	957	8.6	(7.9, 9.2)	1,662	13.1	(12.1, 14.2)
Strongly agree	167	2.4	(2.0, 2.8)	335	3.0	(2.6, 3.4)	730	5.8	(5.0, 6.5)
Q5.4 Harm my career or career prospects									
Strongly disagree	1,730	24.7	(23.6, 25.7)	2,817	25.2	(24.2, 26.1)	3,133	24.8	(23.4, 26.2)
Disagree	1,888	26.9	(25.8, 28.0)	3,159	28.2	(27.3, 29.2)	3,044	24.1	(22.7, 25.4)
Uncertain	1,407	20.1	(19.1, 21.0)	2,380	21.3	(20.4, 22.2)	2,984	23.6	(22.2, 25.0)
Agree	1,406	20.0	(19.0, 21.0)	1,905	17.0	(16.2, 17.9)	2,275	18	(16.8, 19.2)
Strongly agree	586	8.3	(7.7, 9.0)	926	8.3	(7.7, 8.9)	1,207	9.55	(8.6, 10.5)

	Officers				Non-commissioned officers				Other ranks			
	N=7,017				N=11,188				N=12,643			
	N	%	95% CI		N	%	95% CI		N	%	95% CI	
Q5.5 People would treat me differently												
Strongly disagree	1,548	22.1	(21.0, 23.1)		2,523	22.6	(21.6, 23.5)		2,949	23.3	(22.0, 24.7)	
Disagree	1,926	27.4	(26.3, 28.5)		3,086	27.6	(26.6, 28.6)		3,106	24.6	(23.2, 26.0)	
Uncertain	1,598	22.8	(21.7, 23.8)		2,596	23.2	(22.3, 24.1)		3,003	23.8	(22.4, 25.1)	
Agree	1,578	22.5	(21.5, 23.5)		2,321	20.7	(19.8, 21.6)		2,548	20.2	(18.9, 21.4)	
Strongly agree	367	5.2	(4.7, 5.8)		662	5.9	(5.4, 6.4)		1,036	8.2	(7.3, 9.1)	
Q5.6 Seen as weak												
Strongly disagree	1,724	24.6	(23.5, 25.6)		2,838	25.4	(24.4, 26.3)		3,063	24.2	(22.9, 25.6)	
Disagree	2,271	32.4	(31.2, 33.5)		3,577	32.0	(31.0, 33.0)		3,457	27.3	(25.9, 28.8)	
Uncertain	1,339	19.1	(18.1, 20.1)		2,149	19.2	(18.3, 20.1)		2,638	20.9	(19.6, 22.2)	
Agree	1,335	19.0	(18.0, 20.0)		1,925	17.2	(16.4, 18.0)		2,329	18.4	(17.2, 19.7)	
Strongly agree	348	5.0	(4.4, 5.5)		699	6.2	(5.7, 6.8)		1,155	9.14	(8.2, 10.1)	
Q5.7 Stop me from being deployed												
Strongly disagree	1,385	19.7	(18.8, 20.7)		2,079	18.6	(17.7, 19.4)		2,440	19.3	(18.0, 20.6)	
Disagree	1,234	17.6	(16.6, 18.5)		1,987	17.8	(16.9, 18.6)		2,107	16.7	(15.5, 17.9)	
Uncertain	1,859	26.5	(25.4, 27.6)		2,895	25.9	(24.9, 26.8)		3,486	27.6	(26.1, 29.0)	
Agree	1,710	24.4	(23.3, 25.4)		2,598	23.2	(22.3, 24.2)		2,717	21.5	(20.2, 22.8)	
Strongly agree	828	11.8	(11.0, 12.6)		1,628	14.6	(13.8, 15.3)		1,894	15	(13.8, 16.1)	

Table B.41: Prevalence of reported stigma and barriers to care in the ADF, by type and Service

	Navy								
	Male			Female			Total		
	N=6,284			N=1,500			N=7,784		
	N	%	95% CI	N	%	95% CI	N	%	95% CI
Q5.1 Proportion that sought help	1,019	16.2	(14.9, 17.5)	403	26.9	(24.1, 29.7)	1,422	18.3	(17.1, 19.5)
Q5.2 Not knowing where to get help									
Strongly disagree	2,728	43.4	(41.6, 45.3)	735	49.0	(45.8, 52.2)	3,463	44.5	(42.9, 46.1)
Disagree	2,530	40.3	(38.4, 42.1)	572	38.1	(35.0, 41.2)	3,102	39.8	(38.3, 41.4)
Uncertain	590	9.4	(8.2, 10.6)	101	6.7	(5.1, 8.4)	691	8.9	(7.9, 9.9)
Agree	358	5.7	(4.8, 6.6)	77	5.1	(3.6, 6.6)	435	5.6	(4.8, 6.4)
Strongly agree	78	1.2	(0.8, 1.7)	15	1.0	(0.4, 1.7)	93	1.2	(0.8, 1.6)
Q5.3 Difficulty getting time off work									
Strongly disagree	2,149	34.2	(32.4, 35.9)	512	34.1	(31.1, 37.2)	2,661	34.2	(32.7, 35.7)
Disagree	2,265	36.0	(34.3, 37.8)	485	32.4	(29.4, 35.3)	2,750	35.3	(33.8, 36.9)
Uncertain	821	13.1	(11.7, 14.4)	232	15.5	(13.1, 17.9)	1,053	13.5	(12.4, 14.7)
Agree	756	12.0	(10.8, 13.3)	207	13.8	(11.5, 16.0)	963	12.4	(11.2, 13.5)
Strongly agree	294	4.7	(3.8, 5.5)	63	4.2	(2.9, 5.5)	357	4.6	(3.9, 5.3)
Q5.4 Harm my career or career prospects									
Strongly disagree	1,579	25.1	(23.5, 26.7)	361	24.1	(21.3, 26.8)	1,940	24.9	(23.5, 26.3)
Disagree	1,599	25.4	(23.8, 27.1)	373	24.9	(22.1, 27.6)	1,972	25.3	(23.9, 26.7)
Uncertain	1,285	20.4	(18.9, 22.0)	315	21.0	(18.3, 23.6)	1,600	20.5	(19.2, 21.9)
Agree	1,224	19.5	(18.0, 21.0)	299	19.9	(17.4, 22.5)	1,523	19.6	(18.3, 20.9)
Strongly agree	597	9.5	(8.4, 10.6)	153	10.2	(8.2, 12.2)	750	9.6	(8.7, 10.6)

	Army									Air Force								
	Male			Female			Total			Male			Female			Total		
	N=13,692			N=1,834			N=15,526			N=6,193			N=1,345			N=7,538		
	N	%	95% CI	N	%	95% CI	N	%	95% CI	N	%	95% CI	N	%	95% CI	N	%	95% CI
	2,183	15.9	(15.0, 16.9)	508	27.7	(25.4, 30)	2,692	17.3	(16.4, 18.2)	988	16	(14.9, 17.0)	420	31.2	(28.7, 33.7)	1,408	18.7	(17.7, 19.7)
	6,719	49.1	(47.7, 50.4)	1,027	56.0	(53.5, 58.5)	7,746	49.9	(48.6, 51.1)	2,481	40.1	(38.6, 41.5)	688	51.2	(48.4, 54.0)	3,169	42	(40.8, 43.3)
	5,123	37.4	(36.1, 38.8)	611	33.3	(30.9, 35.8)	5,734	36.9	(35.7, 38.1)	2,760	44.6	(43.1, 46.0)	528	39.3	(36.5, 42.0)	3,288	43.6	(42.3, 44.9)
	1,050	7.7	(6.9, 8.5)	107	5.8	(4.5, 7.2)	1,157	7.5	(6.7, 8.2)	500	8.1	(7.3, 8.9)	55	4.1	(3.0, 5.2)	555	7.4	(6.7, 8.1)
	631	4.6	(4.0, 5.2)	70	3.8	(2.8, 4.9)	701	4.5	(4.0, 5.1)	390	6.3	(5.6, 7.0)	66	4.9	(3.7, 6.2)	456	6.1	(5.4, 6.7)
	170	1.2	(0.9, 1.5)	19	1.0	(0.5, 1.5)	189	1.2	(0.9, 1.5)	63	1.0	(0.7, 1.3)	7	0.5	(0.1, 0.9)	70	0.9	(0.7, 1.2)
	5,186	37.9	(36.6, 39.2)	701	38.2	(35.8, 40.6)	5,886	37.9	(36.7, 39.1)	2,306	37.2	(35.8, 38.6)	543	40.4	(37.6, 43.1)	2,849	37.8	(36.5, 39.0)
	4,639	33.9	(32.6, 35.2)	635	34.6	(32.2, 37.1)	5,274	34.0	(32.8, 35.1)	2,462	39.8	(38.3, 41.2)	475	35.3	(32.7, 38.0)	2,937	39.0	(37.7, 40.2)
	1,813	13.2	(12.3, 14.2)	253	13.8	(12.0, 15.6)	2,067	13.3	(12.4, 14.2)	677	10.9	(10.0, 11.9)	165	12.3	(10.5, 14.1)	842	11.2	(10.3, 12.0)
	1,441	10.5	(9.6, 11.4)	189	10.3	(8.7, 12.0)	1,630	10.5	(9.7, 11.3)	576	9.3	(8.5, 10.2)	127	9.4	(7.8, 11.0)	703	9.3	(8.6, 10.1)
	613	4.5	(3.9, 5.1)	55	3.0	(2.1, 3.9)	669	4.3	(3.8, 4.9)	172	2.8	(2.3, 3.3)	35	2.6	(1.7, 3.5)	207	2.7	(2.3, 3.2)
	3,468	25.3	(24.1, 26.5)	471	25.7	(23.4, 27.9)	3,939	25.4	(24.3, 26.5)	1,430	23.1	(21.9, 24.3)	371	27.6	(25.1, 30.1)	1,801	23.9	(22.8, 25.0)
	3,414	24.9	(23.7, 26.1)	523	28.5	(26.2, 30.8)	3,937	25.4	(24.3, 26.4)	1,835	29.6	(28.3, 31.0)	348	25.9	(23.5, 28.3)	2,183	29.0	(27.8, 30.1)
	3,061	22.4	(21.2, 23.5)	350	19.1	(17.1, 21.1)	3,411	22.0	(20.9, 23.0)	1,468	23.7	(22.4, 24.9)	293	21.8	(19.5, 24.1)	1,761	23.4	(22.2, 24.5)
	2,516	18.4	(17.3, 19.4)	324	17.7	(15.7, 19.7)	2,841	18.3	(17.3, 19.3)	998	16.1	(15.1, 17.2)	224	16.7	(14.6, 18.7)	1,223	16.2	(15.3, 17.2)
	1,233	9.0	(8.2, 10.0)	165	9.0	(7.5, 10.5)	1,399	9.0	(8.3, 9.7)	462	7.5	(6.7, 8.2)	108	8.0	(6.5, 9.6)	570	7.6	(6.9, 8.3)

	Navy								
	Male			Female			Total		
	N=6,284			N=1,500			N=7,784		
	N	%	95% CI	N	%	95% CI	N	%	95% CI
Q5.5 People would treat me differently									
Strongly disagree	1,462	23.3	(21.7, 24.8)	330	22.0	(19.3, 24.6)	1,792	23.0	(21.7, 24.4)
Disagree	1,646	26.2	(24.6, 27.8)	380	25.3	(22.6, 28.0)	2,026	26.0	(24.6, 27.4)
Uncertain	1,464	23.3	(21.7, 24.9)	334	22.3	(19.5, 25.0)	1,798	23.1	(21.7, 24.5)
Agree	1,311	20.9	(19.4, 22.4)	340	22.7	(20.0, 25.4)	1,652	21.2	(19.9, 22.5)
Strongly agree	401	6.4	(5.4, 7.3)	116	7.7	(5.9, 9.5)	517	6.6	(5.8, 7.5)
Q5.6 Seen as weak									
Strongly disagree	1,568	25.0	(23.3, 26.6)	392	26.1	(23.3, 28.9)	1,960	25.2	(23.8, 26.6)
Disagree	1,872	29.8	(28.1, 31.5)	418	27.9	(25.0, 30.7)	2,290	29.4	(27.9, 30.9)
Uncertain	1,297	20.6	(19.1, 22.2)	299	20.0	(17.3, 22.6)	1,596	20.5	(19.2, 21.8)
Agree	1,164	18.5	(17.1, 20.0)	279	18.6	(16.1, 21.1)	1,443	18.5	(17.3, 19.8)
Strongly agree	383	6.1	(5.2, 7.0)	112	7.5	(5.7, 9.2)	495	6.4	(5.5, 7.2)
Q5.7 Stop me from being deployed									
Strongly disagree	1,215	19.3	(17.9, 20.8)	315	21.0	(18.4, 23.6)	1,530	19.7	(18.4, 20.9)
Disagree	1,202	19.1	(17.7, 20.6)	229	15.3	(13.0, 17.5)	1,431	18.4	(17.1, 19.6)
Uncertain	1,591	25.3	(23.7, 26.9)	374	24.9	(22.1, 27.7)	1,965	25.2	(23.8, 26.7)
Agree	1,451	23.1	(21.5, 24.6)	376	25.1	(22.4, 27.8)	1,827	23.5	(22.1, 24.8)
Strongly agree	825	13.1	(11.9, 14.4)	206	13.7	(11.5, 16.0)	1,031	13.3	(12.1, 14.4)

	Army									Air Force								
	Male			Female			Total			Male			Female			Total		
	N=13,692			N=1,834			N=15,526			N=6,193			N=1,345			N=7,538		
	N	%	95% CI	N	%	95% CI	N	%	95% CI	N	%	95% CI	N	%	95% CI	N	%	95% CI
	3,196	23.3	(22.2, 24.5)	456	24.9	(22.6, 27.1)	3,652	23.5	(22.5, 24.6)	1,242	20.1	(18.9, 21.2)	333	24.8	(22.4, 27.2)	1,576	20.9	(19.8, 22.0)
	3,449	25.2	(24.0, 26.4)	493	26.9	(24.6, 29.1)	3,941	25.4	(24.3, 26.5)	1,764	28.5	(27.2, 29.8)	387	28.8	(26.3, 31.3)	2,151	28.5	(27.4, 29.7)
	3,145	23.0	(21.8, 24.1)	375	20.4	(18.4, 22.5)	3,520	22.7	(21.6, 23.7)	1,588	25.6	(24.4, 26.9)	292	21.7	(19.4, 24.0)	1,880	24.9	(23.8, 26.1)
	2,885	21.1	(20.0, 22.2)	411	22.4	(20.3, 24.5)	3,296	21.2	(20.2, 22.2)	1,254	20.2	(19.1, 21.4)	246	18.3	(16.2, 20.5)	1,500	19.9	(18.9, 20.9)
	1,018	7.4	(6.7, 8.2)	100	5.4	(4.2, 6.6)	1,117	7.2	(6.5, 7.9)	345	5.6	(4.9, 6.3)	86	6.4	(5.0, 7.8)	431	5.7	(5.1, 6.3)
	3,397	24.8	(23.6, 26.0)	473	25.8	(23.5, 28.1)	3,870	24.9	(23.8, 26.0)	1,444	23.3	(22.1, 24.6)	352	26	(23.7, 28.6)	1,795	23.8	(22.7, 24.9)
	3,940	28.8	(27.5, 30.0)	527	28.8	(26.4, 31.1)	4,468	28.8	(27.6, 29.9)	2,100	33.9	(32.5, 35.3)	448	33	(30.7, 35.9)	2,548	33.8	(32.6, 35.0)
	2,635	19.2	(18.1, 20.3)	339	18.5	(16.5, 20.5)	2,974	19.2	(18.2, 20.2)	1,324	21.4	(20.2, 22.6)	233	17	(15.2, 19.4)	1,557	20.7	(19.6, 21.7)
	2,566	18.7	(17.7, 19.8)	379	20.7	(18.5, 22.8)	2,945	19.0	(18.0, 19.9)	987	15.9	(14.9, 17.0)	215	16	(14.0, 17.9)	1,201	15.9	(15.0, 16.9)
	1,155	8.4	(7.6, 9.2)	116	6.3	(5.1, 7.6)	1,270	8.2	(7.5, 8.9)	339	5.5	(4.8, 6.1)	98	7	(5.8, 8.8)	437	5.8	(5.2, 6.4)
	2,598	19.0	(17.9, 20.1)	397	21.7	(19.6, 23.8)	2,996	19.3	(18.3, 20.3)	1,075	17.4	(16.3, 18.5)	303	22.5	(20.2, 24.8)	1,378	18.3	(17.3, 19.3)
	2,192	16.0	(15.0, 17.0)	331	18.0	(16.1, 20.0)	2,523	16.2	(15.3, 17.2)	1,124	18.1	(17.0, 19.3)	251	18.7	(16.5, 20.8)	1,375	18.2	(17.2, 19.2)
	3,381	24.7	(23.5, 25.9)	424	23.1	(21.0, 25.3)	3,805	24.5	(23.4, 25.6)	2,099	33.9	(32.5, 35.3)	371	27.6	(25.1, 30.1)	2,470	32.8	(31.5, 34.0)
	3,261	23.8	(22.7, 25.0)	419	22.9	(20.7, 25.0)	3,680	23.7	(22.6, 24.8)	1,262	20.4	(19.2, 21.6)	256	19.0	(16.9, 21.2)	1,518	20.1	(19.1, 21.2)
	2,260	16.5	(15.5, 17.5)	262	14.3	(12.5, 16.1)	2,522	16.2	(15.3, 17.2)	633	10.2	(9.3, 11.1)	164	12.2	(10.4, 14.0)	797	10.6	(9.8, 11.4)

Table B.42: Prevalence of reported stigma and barriers to care in the ADF, by type and deployment history

	Never deployed			Deployed		
	N=16,966			N=12,899		
	N	%	95% CI	N	%	95% CI
Q5.1 Proportion that sought help	3,277	17.9	(17.1, 18.7)	2,246	17.9	(17.0, 18.8)
Q5.2 Not knowing where to get help						
Strongly disagree	8,286	45.3	(44.2, 46.3)	6,092	48.6	(47.4, 49.8)
Disagree	7,274	39.7	(38.7, 40.8)	4,850	38.7	(37.5, 39.9)
Uncertain	1,565	8.5	(7.9, 9.2)	838	6.7	(6.0, 7.4)
Agree	979	5.3	(4.8, 5.8)	613	4.9	(4.3, 5.4)
Strongly agree	208	1.1	(0.9, 1.4)	144	1.1	(0.9, 1.4)
Q5.3 Difficulty getting time off work						
Strongly disagree	6,493	35.5	(34.4, 36.5)	4,903	39.1	(37.9, 40.3)
Disagree	6,484	35.4	(34.4, 36.4)	4,476	35.7	(34.5, 36.9)
Uncertain	2,498	13.6	(12.9, 14.4)	1,464	11.7	(10.8, 12.5)
Agree	2,073	11.3	(10.6, 12.0)	1,223	9.8	(9.0, 10.5)
Strongly agree	763	4.2	(3.7, 4.6)	470	3.7	(3.2, 4.3)
Q5.4 Harm my career or career prospects						
Strongly disagree	4,522	24.7	(23.8, 25.6)	3,157	24.1	(24.1, 26.3)
Disagree	4,883	26.7	(25.7, 27.6)	3,208	24.5	(24.5, 26.7)
Uncertain	4,154	22.7	(21.8, 23.6)	2,618	19.9	(19.9, 21.9)
Agree	3,180	17.4	(16.5, 18.2)	2,407	18.2	(18.2, 20.2)
Strongly agree	1,572	8.6	(8.0, 9.2)	1,147	8.4	(8.4, 9.9)
Q5.5 People would treat me differently						
Strongly disagree	4,102	22.4	(21.5,23.3)	2,917	23.3	(22.2,24.3)
Disagree	4,960	27.1	(26.1,28.0)	3,159	25.2	(24.1,26.3)
Uncertain	4,257	23.2	(22.3,24.2)	2,941	23.5	(22.4,24.5)
Agree	3,747	20.5	(19.6,21.3)	2,700	21.5	(20.5,22.6)
Strongly agree	1,245	6.8	(6.2,7.4)	820	6.5	(5.9,7.2)
Q5.6 Seen as weak						
Strongly disagree	4,445	24.3	(23.3,25.2)	3,180	25.4	(24.3,26.4)
Disagree	5,724	31.3	(30.3,32.3)	3,581	28.6	(27.5,29.7)
Uncertain	3,645	19.9	(19.0,20.8)	2,482	19.8	(18.8,20.8)
Agree	3,214	17.6	(16.7,18.4)	2,375	18.9	(18.0,19.9)
Strongly agree	1,283	7.0	(6.4,7.6)	919	7.3	(6.6,8.0)

	Never deployed			Deployed		
	N=16,966			N=12,899		
	N	%	95% CI	N	%	95% CI
Q5.7 Stop me from being deployed						
Strongly disagree	3,526	19.3	(18.4,20.1)	2,377	19.0	(18.0,19.9)
Disagree	3,294	18.0	(17.2,18.8)	2,035	16.2	(15.3,17.1)
Uncertain	5,239	28.6	(27.6,29.6)	3,001	23.9	(22.9,25.0)
Agree	3,863	21.1	(20.2,22.0)	3,162	25.2	(24.1,26.3)
Strongly agree	2,389	13.0	(12.3,13.8)	1,962	15.6	(14.7,16.6)

Annex A data tables

Table B.43: Demographic profile of Phase 1 survey respondents and non-respondents

Characteristic	Population (N=50,049)		Respondents (N=24,481, 48.9%)		Non-respondents (N=25,568, 51.1%)	
	N	%	N	%	N	%
Sex						
Female	6,808	13.6	3,888	15.9	2,920	11.4
Male	43,241	86.4	20,593	84.1	22,648	88.6
Service						
Navy	11,612	23.2	5,392	22.0	6,220	24.3
Females	2,104	4.2	1,053	4.3	1,051	4.1
Males	9,508	19.0	4,339	17.7	5,169	20.2
Army	25,356	50.7	11,429	46.7	13,927	54.5
Females	2,513	5.0	1,437	5.9	1,076	4.2
Males	22,843	45.6	9,992	40.8	12,851	50.3
Air Force	13,081	26.1	7,660	31.3	5,421	21.2
Females	2,191	4.4	1,398	5.7	793	3.1
Males	10,890	21.8	6,262	25.6	4,628	18.1
Age	33.2 (M)	9.2 (SD)	35.5 (M)	9.3 (SD)	30.9 (M)	8.4 (SD)
18–27	18,422	36.8	6,514	26.6	11,908	46.6
28–37	16,688	33.3	8,285	33.8	8,403	32.9
38–47	10,984	21.9	6,899	28.2	4,085	16.0
48–57	3,748	7.5	2,640	10.8	11,008	4.3
58–over	207	0.4	143	0.6	64	0.3
Marital status						
Married	31,500	62.9	18,882	77.1	12,618	49.4
Not married	18,549	37.1	5,599	22.9	12,950	50.6
Education ^a		% (95% CI)				
Missing	–	–	396	1.6		
Primary school	89	0.2 (0.1, 0.2)	47	0.2		
Secondary school up to Grade 10	5,389	10.8 (10.4, 11.1)	2,445	10.0		
Secondary school up to Grade 11–12	15,620	31.2 (30.7, 31.7)	6,831	27.9		
Certificate	11,927	23.8 (23.4, 24.3)	5,268	21.5		
Diploma	6,569	13.1 (12.8, 13.4)	3,487	14.2		
Bachelor degree	5,132	10.3 (10.0, 10.5)	2,888	11.8		

Characteristic	Population (N=50,049)		Respondents (N=24,481, 48.9%)		Non-respondents (N=25,568, 51.1%)	
	N	%	N	%	N	%
Postgraduate	5,322	10.6 (10.4, 10.8)	3,119	12.7		
Length of service (years) ^{a, b}	Mean (95% CI)					
Regular	11.6 (11.5, 11.7)	8.8 (SD)	12.9 (M)	8.9 (SD)		
Reserve	4.4 (4.3, 4.6)	5.0 (SD)	4.6 (M)	5.2 (SD)		
Rank						
Commissioned officer	12,034	24.0	7,268	29.7	4,766	18.6
Non-commissioned officer	22,319	44.6	12,381	50.6	9,938	38.9
Other ranks	15,696	31.4	4,832	19.7	10,864	42.5
MEC status						
MEC 1	32,816	65.6	14,954	61.1	17,862	69.9
MEC 2	11,712	23.4	6,726	27.5	4,986	19.5
MEC 3	4,485	8.9	2,301	9.4	2,184	8.5
MEC 4	1,036	2.1	500	2.0	536	2.1
ADF deployment						
Missing	983	2.0	0	0.0	983	3.8
Yes	32,080	64.1	15,952	65.2	16,128	63.1
No	16,986	33.9	8,529	34.8	8,457	33.1
Months deployed over last 3 years ^a	3.7 (3.7, 3.8)	4.6 (SD)	3.6 (M)	4.5 (SD)		
Months deployed over last 3 years on ship (non-MEAO Navy) ^c	10.6 (10.3, 10.9)	9.7 (SD)	9.8 (M)	9.5 (SD)		
Currently on operational deployment (non-MEAO) ^d		% (95% CI)				
Missing	–	–	252	1.7		
Yes	2,453	8.0 (7.5, 8.4)	1,065	7.4		
No	28,395	92.0 (91.6, 92.5)	13,128	90.9		
Length intending to stay in military (non-MEAO) ^d	11.9 (11.7, 12.0)	10.2 (SD)	12.3 (M)	10.4 (SD)		
Length of service in the ADF (years)	11.6 (M)	8.8 (SD)	13.7 (M)	9.3 (SD)	9.7 (M)	7.8 (SD)

a Population prevalences estimated from 23,156 survey responses.

b Length of service among those who have served as a regular or reservist.

c Population prevalence estimated from 3,156 survey responses.

d Population prevalences estimated from 13,265 survey responses.

Table B.44: Consent and CIDI completion rates

	Total sample	
	No.	%
Consent to complete a survey questionnaire	26,281	N/A
Consent to being contacted to do a telephone interview/for follow-up studies	20,198	76.9
Consent to allow linkage of information contained in my electronic ADF psychological screening records with the study data	21,768	82.8
Consent to allow CMVH to obtain from the ADF, contact details to invite your partner/spouse to participate in a family study	17,554	66.8
CIDI completed	1,798	6.8

Table B.45: Number of days between completing Phase 1 questionnaire and Phase 2 CIDI interview

	Total	28 or less days from completing questionnaire to interview		60 or less days from completing questionnaire to interview		More than 60 days from completing questionnaire to interview	
		No.	%	No.	%	No.	%
CIDI	1,798	640	35.6	1,573	87.5	225	12.5

Table B.46: Demographic profile of the CIDI sample

Characteristic	CIDI sample (N=3,688, 100%)		CIDI respondents (N=1,798, 48.8%)		CIDI non-respondents (N=1,890, 51.2%)	
	No.	%	No.	%	No.	%
Sex						
Females	907	24.6	438	24.4	469	24.8
Males	2,781	75.4	1,360	75.6	1,421	75.2
Service						
Navy	837	22.7	384	21.4	453	24.0
Females	227	6.2	100	5.6	127	6.7
Males	610	16.5	284	15.8	326	17.2
Army	1,325	35.9	716	39.8	609	32.2
Females	322	8.7	165	9.2	157	8.3
Males	1,003	27.2	551	30.6	452	23.9
Air Force	1,526	41.4	698	38.8	828	43.8
Females	358	9.7	173	9.6	185	9.8
Males	1,168	31.7	525	29.2	643	34.0
Age	37.3 (M)	9.4 (SD)	38.3 (M)	9.4 (SD)	36.4 (M)	9.3 (SD)
18–27	733	19.9	300	16.7	433	22.9
28–37	1,183	32.1	546	30.4	637	33.7
38–47	1,246	33.8	662	36.8	584	30.9
48–57	500	13.6	271	15.1	229	12.1
58–over	26	0.7	19	1.1	7	0.4

Characteristic	CIDI sample (N=3,688, 100%)		CIDI respondents (N=1,798, 48.8%)		CIDI non-respondents (N=1,890, 51.2%)	
	No.	%	No.	%	No.	%
Marital status						
Yes	2,862	77.6	1,388	77.2	1,474	78.0
No	826	22.4	410	22.8	416	22.0
Rank						
Officer	1,233	33.4	655	36.4	578	30.6
Non-commissioned officer	1,881	51.0	889	49.4	992	52.5
Other ranks	574	15.6	254	14.1	320	16.9
MEC status						
MEC 1	1,989	53.9	906	50.4	1,083	57.3
MEC 2	1,184	32.1	611	34.0	573	30.3
MEC 3	413	11.2	224	12.5	189	10.0
MEC 4	102	2.8	57	3.2	45	2.4
ADF deployment						
Yes	2,288	62.0	1,111	61.8	1,177	62.3
No	1,400	38.0	687	38.2	713	37.7
Length of service in the ADF	15.3 (M)	9.5 (SD)	16.2 (M)	9.8 (SD)	14.3 (M)	9.2 (SD)

Table B.47: Phase 2: Breakdown of CIDI completers by band and Service

	Navy			Army			Air Force		
	Selected	Completed		Selected	Completed		Selected	Completed	
		No.	%		No.	%		No.	%
Band 3	312	154	49.4	761	433	56.9	366	201	54.9
Band 2	221	98	44.3	279	138	49.5	405	171	42.2
Band 1	304	132	43.4	285	145	50.9	755	326	43.2